



Designation: E 2122 – 04<sub>2</sub>

## Standard Guide for Conducting In-situ Field Bioassays With ~~Marine, Estuarine,~~ and Freshwater Caged Bivalves<sup>1</sup>

This standard is issued under the fixed designation E 2122; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon ( $\epsilon$ ) indicates an editorial change since the last revision or reapproval.

### 1. Scope

1.1 This guide describes procedures for conducting controlled experiments with caged bivalves under field conditions. The purpose of this approach is to facilitate the simultaneous collection of field data to help characterize chemical exposure and associated biological effects in the same organism under environmentally realistic conditions. This approach of characterizing exposure and effects is consistent with the US EPA ecological risk assessment paradigm. Bivalves are useful test organisms for in-situ field bioassays because they (1) concentrate and integrate chemicals in their tissues and have a more limited ability to metabolize most chemicals than other species, (2) exhibit measurable sublethal effects associated with exposure to those chemicals, (3) provide paired tissue chemistry and response data which can be extrapolated to other species and trophic levels, (4) provide tissue chemistry data which can be used to estimate chemical exposure from water or sediment, and (5) facilitate controlled

---

<sup>1</sup> This guide is under the jurisdiction of ASTM Committee E47 on Biological Effects and Environmental Fate and is the direct responsibility of Subcommittee E47.08 on Biological Field Testing.

Current edition approved ~~January~~ May 10, 2004<sub>2</sub>. Published ~~M~~ September 2002. Originally published as E 2122 – 01. Last previous edition E 2122 – 01.

experimentation in the field with large sample sizes because they are easy to collect, cage, and measure **(1, 2)**<sup>2</sup>. The experimental control afforded by this approach can be used to place a large number of animals of a known size distribution in specific areas of concern to quantify exposure and effects over space and time within a clearly defined exposure period. Chemical exposure can be estimated by measuring the concentration of chemicals in water, sediment, or bivalve tissues, and effects can be estimated with survival, growth, and other sublethal end points **(3)**. Although a number of assessments have been conducted using bivalves to characterize exposure by measuring tissue chemistry or associated biological effects, relatively few assessments have been conducted to characterize both exposure and biological effects simultaneously **(2, 4, 5)**. This guide is specifically designed to help minimize the variability in tissue chemistry and response measurements by using a practical uniform size range and compartmentalized cages for multiple measurements on the same individuals.

1.2 The test is referred to as a field bioassay because it is conducted in the field and because it includes an element of relative chemical potency to satisfy the bioassay definition. Relative potency is established by comparing tissue concentrations with effects levels for various chemicals with toxicity and bioaccumulation end points **(6, 7, 8, 9, 10)** even though there may be more uncertainty associated with effects measurements in field studies. Various pathways of exposure can be evaluated because filter-feeding and deposit-feeding are the primary feeding strategies for bivalves. Filter-feeding bivalves may be best suited to evaluate the bioavailability and associated effects of chemicals in the water column (that is, dissolved and suspended particulates); deposit-feeding bivalves may be best suited to evaluate chemicals associated with sediments **(11, 12)**. It may be difficult to demonstrate pathways of exposure under field conditions, particularly since filter-feeding bivalves can ingest suspended sediment and facultative deposit-feeding bivalves can switch between filter- and deposit feeding over relatively small temporal scales. Filter-feeding bivalves caged within 1 m of bottom sediment have also been used effectively in sediment assessments from depths of 10 to 650 m **(5, 13, 14)**. Caged bivalve studies have also been conducted in the intertidal zone **(15)**. The field testing procedures described here are useful for testing most bivalves although modifications may be necessary for a particular species.

1.3 These field testing procedures with caged bivalves are applicable to the environmental evaluation of water and sediment in marine, estuarine, and freshwater environments with almost any combination of chemicals, and methods are being developed to help interpret the environmental significance of accumulated chemicals **(6, 7, 9, 16, 17)**. These procedures could be regarded as a guide to an exposure system to assess chemical bioavailability and toxicity under natural, site- specific conditions, where any clinical measurements are possible.

1.4 Tissue chemistry results from short- and long-term exposures can be reported in terms of concentrations of chemicals in bivalve tissues (for example,  $\mu\text{g/g}$ ), amount (that is, weight or mass) of chemical per animal (for example,  $\mu\text{g/animal}$ ), rate of uptake, or bioaccumulation factor (BAF, the ratio between the concentration of a chemical in bivalve tissues and the concentration in the external environment, including water, sediment, and food). Tissue chemistry results can only be used to calculate a BAF because caged bivalves in the field are exposed to multiple sources of chemicals and can accumulate chemicals from water, sediment, and food. Toxicity results can be reported in terms of survival **(3, 18)**, growth rate **(3, 18)**, or reproductive effects **(19, 20)** after a defined exposure period.

1.5 Other modifications of these procedures might be justified by special needs or circumstances. Although using appropriate procedures is more important than following prescribed procedures, results of tests conducted using unusual procedures are not likely to be comparable to results of standardized tests. Comparisons of results obtained using modified and unmodified versions of these procedures might provide useful information concerning new concepts and procedures for conducting field bioassays with bivalves.

1.6 This guide is arranged as follows:

	Section
Referenced Documents	2
Terminology	3
Summary of Guide	4
Significance and Use	5
Interferences	6
Hazards	7
Experimental Design	8
Apparatus	9
Facilities	
Construction Materials	
Cages	
Test Organisms	10
Species	
Commonly Used Taxa	
Size and Age of Test Organisms	
Source	
Number of Specimens	
Collection	
Handling	
Holding	

<sup>2</sup> The boldface numbers in parentheses refer to references at the end of this standard.

Animal Quality	
Field Procedures	11
Test Initiation: Presort	
Final Measurements and Distribution	
Attachment of PVC Frames	
Deployment	
Retrieval and End-of-Test Measurements	
Analysis of Tissues for Background Contamination	
Collection and Preparation of Tissues for Analysis	
Quality Assurance/Quality Control Procedures	
Sample Containers, Handling, and Preservation	
Ancillary Methodology	12
Temperature	
Food	
Acceptability of Test	13
Report	14
Keywords	15
References	

1.7 The values stated in the International System of Units (SI) (the Modernized Metric System) are to be regarded as standard.

1.8 *This standard may involve hazardous materials, operations, and equipment – particularly during field operations in turbulent waters or extreme weather conditions. This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory requirements prior to use.* Specific hazard statements are given in Section 7.

## 2. Referenced Documents

### 2.1 ASTM Standards:

- D 1129 Terminology Relating to Water<sup>3</sup>
- D 3976 Practice for Preparation of Sediment Samples for Chemical Analysis<sup>4</sup>
- D 4447 Guide for Disposal of Laboratory Chemicals and Samples<sup>5</sup>
- E 380 Practice for Use of the International System of Units (SI) (the Modernized Metric System)<sup>6</sup>
- E 724 Guide for Conducting Static Acute Toxicity Tests Starting with Embryos of Four Species of Saltwater Bivalve Molluscs<sup>7</sup>
- E 729 Guide for Conducting Acute Toxicity Tests on Test Materials with Fishes, Macroinvertebrates, and Amphibians<sup>7</sup>
- E 943 Terminology Relating to Biological Effects and Environmental Fate<sup>7</sup>
- E 1022 Practice for Conducting Bioconcentration Tests with Fishes and Saltwater Bivalve Molluscs<sup>7</sup>
- E 1023 Guide for Assessing the Hazard of a Material to Aquatic Organisms and Their Uses<sup>7</sup>
- E 1191 Guide for Conducting Life-Cycle Toxicity Tests with Saltwater Mysids<sup>7</sup>
- E 1192 Guide for Conducting Acute Toxicity Test on Aqueous Ambient Samples and Effluents with Fishes, Macroinvertebrates, and Amphibians<sup>7</sup>
- E 1342 Practice for Preservation by Freezing, Freeze-Drying, and Low Temperature Maintenance of Bacteria, Fungi, Protista, Viruses, Genetic Elements, and Animal and Plant Tissues<sup>7</sup>
- E 1367 Guide for Conducting 10-day Static Sediment Toxicity Tests with Marine and Estuarine Amphipods<sup>7</sup>
- E 1391 Guide for Collection, Storage, Characterization, and Manipulation of Sediments for Toxicological Testing<sup>7</sup>
- E 1525 Guide for Designing Biological Tests with Sediments<sup>7</sup>
- E 1688 Guide for Determination of the Bioaccumulation of Sediment-Associated Contaminants by Benthic Invertebrates<sup>7</sup>
- E 1706 Test Methods for Measuring the Toxicity of Sediment-Associated Contaminants with Fresh Water Invertebrates<sup>7</sup>
- E 1847 Practice for Statistical Analysis of Toxicity Tests Conducted Under ASTM Guidelines<sup>7</sup>

## 3. Terminology

### 3.1 Definitions:

3.1.1 The words “must,” “should,” “may,” “can,” and “might,” have very specific meanings in this guide. “Must” is used to express an absolute requirement, that is, to state that a test ought to be designed to satisfy the specified condition, unless the purpose of the test requires a different design. “Must” is only used in connection with factors that directly relate to the acceptability of the test. “Should” is used to state that a specified condition is recommended and ought to be met if possible. Although violation of one “should” is rarely a serious matter, violation of several will often render the results questionable. Terms such as “is desirable” are used in connection with less important factors. “May” is used to mean “is (are) allowed to,” “can” is used to mean “is (are) able to,” and “might” is used to mean “could possibly.” Thus the classic distinction between “may” and “can” is preserved and “might” is never used as a synonym for either “may” or “can.”

<sup>3</sup> Annual Book of ASTM Standards, Vol 11.01.

<sup>4</sup> Annual Book of ASTM Standards, Vol 11.02.

<sup>5</sup> Annual Book of ASTM Standards, Vol 11.04.

<sup>6</sup> Discontinued; Replaced by IEEE/ASTM SI-10.

<sup>7</sup> Annual Book of ASTM Standards, Vol 11.05.

3.1.2 For definitions of other terms used in this guide, refer to Terminology D 1129, Guide E 729, Terminology E 943, and Guide E 1023. For an explanation of units and symbols, refer to Practice E 380.

### 3.2 Definitions of Terms Specific to This Standard:

3.2.1 *bioaccumulation factor (BAF)*—the ratio of tissue chemical residue to chemical concentration in the external environment. BAF is measured at steady state in situations where organisms are exposed from multiple sources (that is, water, sediment, food), unless noted otherwise.

3.2.2 *bioassay*—an experiment that includes both an estimate of toxicity and an estimate of relative potency.

3.2.3 *bioavailability*—the fraction of the total chemical concentration in water, on sediment particles, and on food that is available for bioaccumulation.

3.2.4 *biomonitoring*—use of living organisms as “sensors” in water or sediment quality surveillance to detect changes in an effluent or water body or to indicate whether aquatic life may be endangered.

3.2.5 *chemical concentration*—the ratio of the weight or volume of chemicals to the weight or volume of a test sample.

3.2.6 *chemical content*—mass of chemical per whole animal (for example,  $\mu\text{g}/\text{animal}$ ) can be used to normalize the expression of chemical uptake per unit time by eliminating the effects of growth on changing tissues masses.

3.2.7 *chemical fingerprinting*—the use of specific patterns in the ratios of chemicals accumulated in bivalve tissues to identify chemical sources; for example, the ratio of PAH alkylated homologs to parent compounds.

3.2.8 *compartmentalized cage*—a rigid or flexible mesh cage with individual compartments for holding bivalves in a controlled position so that multiple measurements can be made on the same individual organism. The compartmentalized cage helps maximize water flow around individual test organisms and provides even exposure to the test environment.

3.2.9 *growth dilution*—a process whereby the rate of accumulation is exceeded by the rate of tissue growth so that when the concentration is expressed on mass of chemical per mass of tissue over time, it appears as though depuration or elimination is occurring because the concentration ( $\mu\text{g}/\text{g}$ ) is decreasing.

3.2.10 *reference station*—a station similar to the test station(s) in physical and chemical characteristics and with relatively little to no contamination by the particular chemical(s) under study. A reference station should ideally contain only background concentrations of chemicals characteristic of the region.

3.2.11 *scope for growth*—an integrated physiological measure of the energy status of an organism at a particular time, based on the concept that energy in excess of that required for normal maintenance will be available for the growth and reproduction of the organism.

3.2.12 *shell length*—the distance from the tip of the umbo to the distal valve edge.

3.2.13 *site*—a geographical area within a somewhat defined boundary that is being studied. The size of a site is dependent on the extent of suspected perturbation, generally on the order of 0.1 to 50– $\text{km}^2$ . Part of the vagueness in size is due to variability in spatial scale and inadequate results from preliminary reconnaissance survey that clearly define the boundary of suspected stressors.

3.2.14 *steady state*—the state in which fluxes of material moving bidirectionally across a membrane or boundary between compartments or phases have reached a balance. An equilibrium between the phases is not necessarily achieved.

3.2.15 *station*—a specific sampling location or area within a site. The size of a station can vary from a single point with one cage to an area of approximately 10 by 10 m including several cages. Vagueness in size is due to variability in spatial scale and experimental design. Several stations in a small geographic area could comprise a site.

3.2.16 *tissue loss magnification*—the process whereby the tissue mass is lost during the exposure period and the chemical mass remains constant over time, so that when the concentration is expressed on mass of chemical per mass of tissue over time, it appears as though bioaccumulation is occurring because the concentration ( $\mu\text{g}/\text{g}$ ) is increasing.

3.2.17 *uptake*—acquisition of a substance from the environment by an organism as a result of any active or passive process.

3.2.18 *whole-animal wet-weight*—the wet weight (g) of the entire bivalve, including water trapped between the valves.

## 4. Summary of Guide

4.1 This guide describes procedures for exposing marine, estuarine, and freshwater bivalves to chemicals in water, sediment, or water and sediment combined, food in the field under natural in-situ field conditions. The purpose of this guide is to provide a standard approach for in-situ testing with bivalves. Because of its application to a wide variety of species, many of which have a range of tolerance limits for water quality conditions, it is outside the scope of this guide to provide the tolerance limits for all water quality conditions for all species that can be used for in-situ testing. Tolerance limits ~~will be~~ are provided for selected species as examples and points of reference. (6.4)

4.2 The approach can be used ~~in a variety of applications~~ to characterize exposure and effects over space and time. The primary measurement end points are bioaccumulation of chemicals in bivalve tissues to assess biological availability or bioaccumulation potential, and sublethal effects, like growth, to assess adverse biological effects. The bioavailability of chemical(s) in water, sediment, and sediment food and associated biological effects are determined by the relative differences in these exposure and effects end points among stations over time.

4.3 In practice, the two most commonly measured effects end points are survival and growth. Survival is the easiest effects end point to measure and provides an estimate of toxicity in short- or long-term exposures. The survival end point ~~is relatively~~ may be insensitive for ~~many some~~ chemicals but can provide important corroborative effects information. ~~Growth is a~~ Sublethal

endpoints like growth are generally more sensitive sublethal end point. It sensitive. Growth can be determined estimated from changes in whole-animal wet-weight, shell length, tissue weight, or shell weight, with baseline tissue and shell weights for the entire test population estimated from a subsample of that population. Reproduction is another sensitive effects end point, but is more difficult to measure.

4.4 Bioaccumulation and growth are compared among test stations for ranking purposes, among reference and treatment stations, or among stations for temporal and spatial variability as well as short- and long-term trends. It is also possible to use the data to construct dose-response relationships (6, 7) and to identify sources of point and non-point discharges by comparing bioaccumulation and biological effects at various distances away from suspected sources of contamination in a gradient approach (21).

4.5 It is highly recommended that the concentration of chemicals in water and sediment be determined, in addition to measuring the concentration of chemicals in tissues, to characterize total potential exposure; that is, exposure from both water and sediment. These data can be used to establish relationships between chemicals in various environmental compartments as a form of field validation or to establish first-order approximations of these relationships that are validated in the laboratory. Results of the bivalve field bioassay can be used to predict bioaccumulation and biological effects likely to occur in other aquatic organisms under comparable field conditions. While this may only represent first order approximations in some cases, equilibrium partitioning theory, quantitative structure activity relationships, and critical body residue theory suggest that tissue burdens of chemicals associated with adverse effects may be similar across species (6, 7).

## 5. Significance and Use

5.1 The ecological importance of bivalves, their wide geographic distribution, ease of handling in the laboratory and the field, and their ability to filter and ingest large volumes of water and sediment particles make them appropriate species for conducting field bioassays to assess bioaccumulation potential and associated biological effects. The test procedures in this guide are intended to provide guidance for conducting controlled experiments with caged bivalves under “natural,” site-specific conditions. It is important to acknowledge that a number of “natural” factors can affect bivalve growth and the accumulation of chemicals in their tissues (Section 6, Interferences). For this reason, it is highly recommended that supplementary measurements be made on those factors most likely to affect bioaccumulation and growth such as temperature, food (see 6.6), particulate matter, turbidity, salinity, dissolved oxygen, pH, and the velocity of water currents. These measurements are more easily made at shallower depths but may become problematic at the greater depths where caged bivalves have been deployed (for example, 650 m). Nevertheless, it is important to measure the chemicals in water or sediment, and this has been accomplished even at these great depths with discrete samples (for example, (13, 14)). In-situ temperature monitoring devices are available to measure temperatures for periods up to 1 year at depths up to 330 m. Standardized procedures should be used to measure chemicals of concern in water and sediment. Other water or sediment parameters that can affect bioaccumulation and growth, such as temperature, salinity, dissolved oxygen, and pH, could also be measured as part of these field bioassays to provide site-specific information. These data can be used to help explain the measured exposure and effect. This field bioassay can also be conducted in conjunction with laboratory bioassays to help answer questions raised in the field exposures. The field exposures can also be used to validate the results of laboratory bioassays.

5.2 The ultimate resources of concern are communities. However, it is often difficult or impossible to adequately assess the ecological fitness or condition of the community or identify and test the most sensitive species. Bivalves are recommended as a surrogate test species for other species and communities for the following reasons: (1) They readily accumulate many chemicals and show sublethal effects associated with exposure to those chemicals (2); (2) they accumulate many chemicals through multiple pathways of exposure, including water, sediment, and food (22, 23, 24, 25, 26, 27), and (3) caged bivalves have been shown to represent effects on the benthos more accurately than traditional laboratory tests (28, 29). Although bivalve species might be considered insensitive because of their wide use as indicators of chemical bioavailability, it has been suggested that sensitivity is related to the type of test, end points being measured, and duration of exposure (2). In short-term toxicity assessments in which survival is the end point, bivalves may appear to be more tolerant to and less affected by chemicals because of their ability to close their valves for short periods and avoid exposure (30, 31, 32, 33). However, studies comparing the mortality end point in bivalves and other test species have found bivalves to be equally (34, 35) or more sensitive (36, 37) than the other species (Table 1). When the bivalve growth end point was compared to the mortality end point in other test species, the bivalve growth end point was more sensitive (18, 28, 29, 38, 39).

5.2.1 Chronic tests designed to monitor sublethal end points, such as growth, are recommended because bivalves generally show increasing sensitivity with increasing exposure period. Sublethal end points measured in bivalves that have demonstrated high levels of sensitivity include growth (3, 18), reproduction (19), DNA damage (40, 41), metallothioneins and other biochemical markers (42, 43, 44).

5.2.2 There are many field monitoring programs in the US which use bivalves, including the NOAA Status and Trends Program (45), the California Mussel Watch (46), and the California Toxics Monitoring Program, a freshwater monitoring program (47). Similar field-monitoring programs exist in other countries. Numerous laboratory studies throughout the world have examined bioaccumulation and biological effects in bivalves. The existing databases which have compiled bioaccumulation and effects in bivalves and other species (8, 9) make it possible to use tissue residues associated with effects in bivalves as surrogates to estimate effects in both water column and benthic organisms in many freshwater, estuarine, and marine environments.

**TABLE 1 Relative Sensitivity of Bivalves Compared to Other Test Species**

Bivalve Species	Species Compared	Exposure	End Point	Sensitivity
<i>Anodonta grandis</i> (35) (giant floater; currently <i>Pyganodon grandis</i> )	daphnia, fathead minnow, rainbow trout	municipal effluent	LC-50	equal
<i>Anodonata imbecilis</i> (36) (paper pondshell; currently <i>Utterbackia imbecilis</i> )	daphnia	pulp and paper mill effluent	10-d vs 7-d mortality	more
<i>Anodonata imbecilis</i> (34) (paper pondshell; currently <i>Utterbackia imbecilis</i> )	daphnia, midge, fathead minnow	metals	7-d mortality	equal
<i>Musculium transversum</i> (37) (fingernail clam)	17 different species	ammonia	20-d mortality	more sensitive than 16 species
<i>Mercenaria mercenaria</i> (28, 29) (hard clam)	2 amphipods, microtox	sediment	7-d growth, 10-d mortality	more
<i>Mullinia lateralis</i> (38) (dwarf surf clam)	Caged <i>Mercenaria</i> amphipod	more sensitive than lab <i>Mercenaria</i> sediment	(28, 29) 7-d growth, 10-d mortality	more
<i>Mytilus galloprovincialis</i> (18) (Mediterranean mussel)	amphipod	in-situ water column	84-d growth, 10-d mortality	more, [tissue TBT]

5.3 Bivalves are an abundant component of many soft bottom marine, estuarine, and freshwater environments. Intertidal marine bivalves make up a significant portion of many habitats and provide habitats for many additional species. It is important to monitor freshwater bivalves for the following reasons: they are among the first taxa to disappear from benthic communities impacted by chemical pollution and chemicals; they have been shown to be more sensitive than several other major taxa in laboratory tests. (48) The threatened and endangered status of many freshwater bivalve species also make them an important group to monitor.

5.4 If practical, the species to be used in a field bioassay should be one that is endemic to the area under investigation. In many cases, the specific area under investigation may not support bivalves due to a variety of factors including high concentrations of chemicals, competition or predation, or lack of suitable habitat or substrate. Under these conditions, it may be desirable to use a species that would normally be found in the environment if all conditions were favorable; however, it may be necessary to use a surrogate species, that is, a species that can tolerate the environmental conditions but is not normally found in the area, if native species are unavailable in the test area.

5.5 ~~With the current invasion of unwanted exotic species such as zebra mussels (*Dreissena polymorpha*) and Asiatic clams (*Corbicula fluminea*) in freshwaters and clams (*Potamocorbula*) in marine waters, it is important to verify that these species are already found in the area of concern before using them for in-situ field bioassays. Zebra mussels, Asiatic clams, and *Potamocorbula* are good candidates for in-situ field bioassays and have been used successfully (for example, (49, 50, 51, 52)), but these species should be used with caution. Planning for regional and local permits sometimes takes considerable periods of time and the planning process should therefore begin early. Other unwanted introductions include parasites or diseases (*Perkinsus marinus*) from infected bivalves such as oysters. Similarly, unwanted pests could be introduced from microscopic attached forms on the shells of transplanted bivalves such as the freshwater weed *Hydrilla hydrilla*. A potential problem that is unique to freshwater bivalves is the introduction of glochidia stages that are parasitic on fish for a portion of their life cycle. Although this is common in areas where the freshwater bivalves naturally reside, potential effects should be considered during the planning of in-situ field bioassays (for example, see 10.3).~~

#### 5.6 Bivalves

5.5 Bivalves generally utilize one of two primary modes of feeding: filter-feeding or deposit feeding. However, all known deposit-feeding bivalves are facultative in that they can either deposit- or filter-feed. Filter-feeders assimilate dissolved organics as well as suspended particulate matter, including plankton and suspended sediments, from the water column and have the potential for exposure to chemicals associated with this ingested material. Facultative deposit-feeding bivalves can be exposed to chemicals associated with sediments as they ingest sediments. They also ingest particulate material from the water column as they filter feed. As such, bivalves are capable of integrating exposure to chemicals dissolved in water and sorbed on sediment particles on the bottom or in suspension. It should be acknowledged that bivalves transplanted in the overlying water above sediment or transplanted directly on or in sediment may not exclusively accumulate or be affected by chemicals in a particular medium. That is, bivalves in or on sediment may still filter and accumulate chemicals from overlying water. Conversely, bivalves transplanted in the water column may filter suspended sediment and accumulate chemicals from that sediment. Bivalves can also assimilate chemicals as they ventilate overlying water.

5.76 Field bioassays are conducted to obtain information concerning the bioavailability of chemicals in the water column or bedded sediments and subsequent biological effects on bivalves after short- and long-term exposure to water and sediment under site-specific conditions. ~~Results of these exposure and effects field studies can be useful for studying biological availability of chemicals, and structure-activity relationships, among stations or groups of stations or sites.~~ These bioassays do not necessarily provide information about whether delayed effects will occur, although a post-exposure observation period could provide such information. Sublethal post-exposure observations may include gonad development, spawning success, gamete survival, and

development. The decision to conduct post-exposure studies in the field or in the laboratory depends on the observations being made, test conditions required, and experimental logistics.

5.8 At a minimum, biological effects should be characterized by measuring survival and growth. Survival may not always be a very sensitive indicator of effects in bivalves (3), but it is an important parameter to monitor. Several factors can affect survival, including handling prior to test initiation and physical-chemical factors at the deployment stations. Survival can be easily quantified, although it is possible to have some individuals missing at the end of the test due to shell decomposition. Under some circumstances, more individuals may be present at the end of the test than at the start. This would most likely be due to the settlement of juvenile bivalves during the course of the test. This can easily be accounted for as new recruits should be smaller than the test bivalves. All recruits should be removed prior to determining survival and assessing effects end points. Bivalve survival should be confirmed before proceeding with other effects measurements. However, it is possible for shells to stick together due to mucilaginous material or sediment within the shells, prohibiting a precise determination of death. Thus, all dead bivalves may not be identified until the tissue removal process when the shells are opened to reveal the internal tissues.

5.9 Growth is a sensitive sublethal effects end point that is easy to measure and is recommended for all field bioassays. It is more sensitive than mortality, and reductions in growth have been related to adverse effects on bivalve populations (1). As many growth end points as are practical should be measured for assessing growth. For example, it has been shown that shell growth and tissue growth are decoupled. Measuring only one of these end points could provide misleading results and lead to a spurious interpretation of environmental effects on growth (53, 54). Growth end points include, but are not limited to, whole-animal wet weight, shell length, tissue weight, shell weight (18). Whole-animal wet-weights and shell lengths are nondestructive measurements and can be made multiple times over the course of the exposure period. At a minimum, whole-animal wet-weights and shell lengths should be measured at the beginning and end of the test. Since tissue weights and shell weights provide a different perspective on animal health and may be related to different stressors, they should also be measured at the beginning and end of the test (18). Because these measurements are destructive, beginning-of-test tissue and shells weights for bivalves to be deployed can be estimated from a subsample of the bivalves distributed to the individual cages. The bivalves used to estimate beginning-of-test tissue and shell weights should be within the same size range as those identified for deployment. Because the initial tissue and shell weights are based on a subsample of the test population, the change in these metrics over the test period is an indirect determination and has some uncertainty. However, tissue and shell weights can provide the most discriminating measurements under certain conditions, particularly when growth rates are slow (18).

5.10 The

5.7 The in-situ exposures described in this guide could be followed by laboratory measurements, such as scope for growth (2), filtration rate (55), byssal thread production (56, 57, 58), and biomarkers (59, 60).

5.11 Although tissue dry weights are less variable than wet weights, drying the tissues has some limitations. (1) It is more time consuming to dry all the tissues and make the weight measurements. (2) In a combined bioaccumulation and biological effects test, the same wet tissues can be used for chemical analysis and wet-weight measurements (drying tissues may destroy organic chemicals, and limit their ability to predict bioaccumulation potential). (3) A wet-weight approach has been used successfully (18), and may provide better correlations with other growth metrics. Nevertheless, if additional testing clearly demonstrates an advantage to measuring dry weights, or if particular studies require more emphasis on the accuracy of tissue weight measurements, it would be relatively simple to alter the procedures accordingly.

5.12 The

5.8 The bivalve field bioassay can be used to determine the spatial or temporal trends of chemical bioavailability in water and sediment and effects due to exposure to those chemicals. Spatial comparisons of parameters of concern can be made by distributing the caged bivalves along physical and chemical gradients at scales commensurate with the desired level of discrimination. For example, station locations might be distributed along a known physical or chemical gradient in relation to the boundary of a disposal site (61, 62, 63, 64, 65), sewage outfall (66), or effluent pipe or at stations identified as containing elevated concentrations of chemicals in water or sediment as identified in a reconnaissance survey (3, 67, 68). This can be accomplished by placing caged bivalves along horizontal transects or at different depths in the water column. Temporal comparisons can be made by conducting before-and- after studies. For example, the effectiveness of dredge activities, effluent diffuser construction, effluent reduction, or remedial action can be determined by conducting field bioassays before the action, during the action, and after the action.

5.139 The relative bioavailability of chemicals from the various pathways of exposure (that is, aqueous phase, suspended particulate matter, sediment) and subsequent effects can be determined by simultaneously deploying bivalves with different feeding strategies and making supplementary measurements. A combination of filtration and the use of sediment traps followed by chemical analysis of the various environmental compartments can be used to identify the relative contribution of the aqueous phase, suspended particulate matter, and sediment. Lipid bags or semi-permeable membrane devices (SPMDs), which predominantly collect the dissolved fraction of chemicals, could also be used ( 69, 70, 71, 72, 73). The bioaccumulation of chemicals and effects among different bivalve species deployed side-by-side can be compared and used to help explain the spatial variability of chemical contamination, particularly if the different species are placed in different locations (that is, in the water column, on top of the sediments, within the sediments). This field assessment approach could be supplemented with laboratory studies designed to answer specific questions regarding dissolved versus particulate pathways of exposure.

5.140 Results of bivalve field bioassays might be an important consideration when assessing the hazards of materials to aquatic

organisms (see Guide E 1023) or when deriving water or sediment quality guidelines for aquatic organisms (15, 74). They might also be useful for ~~determining~~ establishing tissue residue criteria. Bivalve field bioassays can be useful in making decisions regarding the extent of remedial action needed for contaminated sites. They also provide a convenient method for manipulative field experiments, hypothesis testing, and monitoring specific sites before, during, and after cleanup operations (67, 68) .

## 6. Interferences

6.1 As with all bioassay procedures, there are limitations to the methods described in this guide. However, these limitations should not be considered as a reason for not using the methods described in this guide.

6.2 Results of bivalve field bioassays will depend, in part, on natural factors, including temperature, food supply, other physical and chemical properties of the test environment, selection of adequate reference areas, species selected, condition of the test organisms, exposure technique, and handling of the bivalves prior to deployment. Taking bivalves out of their habitat and weighing and measuring them may be stressful to the bivalves. The degree of handling, holding time, and differences between water and sediment conditions at the collection site versus the transplant site may also be stressful. Careful handling and appropriate acclimation can minimize these stresses.

6.3 Condition of the test organisms is critical to the success of the field bioassay. The most important consideration is spawning cycle because of possible interferences on bioaccumulation and growth and with subsequent data interpretation. Generally, chemicals are lost during spawning, resulting in potential underestimation of chemical bioavailability (75). Conversely, the energy used for gonad development and spawning can make bivalves more sensitive to chemicals, reduce their growth rates, and overestimate potential toxicity. Tests should be conducted with populations that will most likely not spawn during the exposure period. The spawning cycle of candidate test species should be evaluated prior to developing the study design, and species that do not spawn during the proposed exposure period should be selected.

6.4 Temperature of the test environment could affect both bioaccumulation and biological effects. Water temperatures should be monitored over the course of the study to quantify the exposure conditions and the potential effects of temperature. As a general guide, examples of temperature tolerance for the most commonly used species are provided in Table 2. Temperature conditions during the exposure period can be quantified using in-situ monitoring devices. These devices can be attached to the deployment cages and set to collect temperature data at specified time intervals for the duration of the test.

6.5 Lack of acclimation to deployment water quality conditions could be an interference. If water quality conditions differ at collection and deployment sites, it may be necessary to acclimate the test organisms gradually to the deployment conditions. This transition is particularly important near the bivalve's tolerance limits and may be accomplished using serial water dilutions until the proper water quality conditions (for example, temperature, salinity, and pH) are reached. Acclimation for temperature should proceed no faster than 3°C in 72 h (Guides E 1022 and E 1688). Once acclimated, bivalves should be maintained under these conditions for a minimum period of time. Holding bivalves for extended periods under laboratory conditions can induce stress because bivalves are particularly sensitive to temperature, nutrition, and water flow. If test specimens are held for an extended period of time in the laboratory, the effect of this holding can be assessed by comparing soft tissue weights, or other indicators of bivalve health, to that of bivalves of the same size group freshly collected from the field. Alternatively, bivalves could be acclimated in the field ~~under reference conditions,~~ conditions similar to the proposed transplant sites.

6.6 Food supply is important because it affects both biological availability and associated biological effects. Food availability may be more difficult to quantify during the test than temperature or other physical factors. Until in-situ monitoring devices for chlorophyll and other nutrient sources are developed, it is suggested that food availability be estimated at least three times during the study (that is, beginning, middle, and end of test). The measurements made (that is, chlorophyll-a, particulate or total organic carbon, and suspended solids) will depend on the feeding strategy of the test species.

6.7 Current speed is important for filter-feeding bivalves because currents regulate the food supply to the test organisms. Currents are also important to facultative deposit-feeding and filter-feeding bivalves in the benthos because flushing may reduce the potential effects of chemicals by dilution with clean water from outside the assessment area. Currents can be quantified during the exposure period with a continuously recording, in-situ current meter or quantified intermittently during the suggested sampling intervals used to measure food availability.

6.8 Salinity is particularly important in estuarine areas, where salinity can range from 0 ppt at the head of a river to 33 ppt at the mouth. Salinity should be evaluated prior to species selection. If there is a wide salinity range, it may be necessary to identify two or more bivalve species for the assessment: one species for the lower end of the salinity range and another for the upper end of the salinity range. It is recommended that both species be deployed in the area where salinity is in the middle, as this provides a means to compare results between species.

6.9 Possible interferences influencing retrieval of test organisms from the field include caged bivalves being washed away during storm events, buried by underwater sediment shifts, theft, vandalism, fouling, disease, and consumption by predators.

6.10 Depending on the environment under assessment, it is possible for the bivalve cages, including the external predator mesh (see 11.3) and the mesh bags, to become fouled with both epiphytic plant and animal growth. Fouling occurs most frequently in highly productive embayments or areas with restricted flow, such as marinas. Excessive fouling can reduce or eliminate flow of water through the cage material, resulting in highly stressful conditions to the test bivalves. If such conditions are anticipated, the deployed cages should be examined for fouling at regular intervals during the exposure period. Fouling organisms can be removed from the exterior surfaces of the cages by hand or with a stiff brush. If the cages are heavily fouled and it is difficult to remove

**TABLE 2 Temperature (°C) and Salinity (Parts per Thousand (ppt)) Tolerance Limits for Selected Bivalve Species  
(Months when spawning may occur and species distribution are also shown)**

Species and Reference	Temperature Range	Salinity Range	Spawning	Distribution
<i>Corbicula fluminea</i> (Asian clam) (76)	2–25	0–5	may be continuous, usually twice/year spring/early summer; later summer	All west, gulf, and east coastal United States to DE River; NM; OH & MS River systems
<i>Dreissena polymorpha</i> (Zebra mussel) (77)	<0–35	0–6	May to September	Canada and Northeastern United States; Great Lakes, St. Lawrence River; MS, OH, IL & TN River drainages; NY Canals, Hudson River, Finger Lakes
<i>Elliptio complanata</i> (Eastern Elliptio) (78, 79, 80)	0–30	0–3	most June to July; some May to September	Gulf St. Lawrence to GA; Great Lakes, except Lake Michigan & Lake Erie
<i>Pyganodon (Anodonta) grandis</i> (floater mussel) (78)	0–30	0–3	most April to May; some to late August	Canada Interior & St. Lawrence River drainage; Hudson Bay, MI and MO Rivers drainages; NM, CO, TX, Mex
<i>Rangia cuneata</i> (Atlantic Rangia) (81, 82)	8–32	0<19	VA: early April to summer; FL: July–November; LA: Mar–May and late summer to November; Mexico: February–June and September to November	Gulf of Mexico coast from northwest FL to Campeche, Mexico; along Atlantic coast to NJ
<i>Argopecten irradians</i> (Bay scallop) (82)	>7 >30	>14–28	mid-Atlantic: mid-April through early September; NY: June and July; NC and FL: August and December	Atlantic coast; Cape Cod to Gulf of Mexico
<i>Crassostrea gigas</i> (Pacific oyster) (82)	4–24	25–35	July to August	Pacific coast; Pacific Northwest
<i>Crassostrea virginica</i> (Eastern oyster) (83)	-2–36	5–32	Gulf of Mexico: April–October; Malpeque Bay, PEI: July–August; Bideford River Estuary, PEI: July	Gulf of Mexico to Cape Cod
<i>Macoma balthica</i> (Baltic clam) (84, 85, 86)	-2–23	5–30	June–August (Europe); July–September (United States)	Greenland to France; Baltic and Wadden Seas; UK; N. Canada to Chesapeake; AK to San Francisco Bay
<i>Mercenaria mercenaria</i> (Hard clam) (87)	<0–35	12–35	March–November depending on latitude and temperature. Peaks in July	Atlantic and Gulf coasts; abundant MA to VA
<i>Mya arenaria</i> (Soft-shell clam) (88)	-1.7–32	10–32	June–September; once/year north of Cape Cod, twice/year south of Cape Cod	Atlantic coast from Labrador to SC; less in FL; AK and CA
<i>Mytilus californianus</i> (California mussel) (82)	7–28	25–33	Continuous throughout year; peaks in July and December	AK to southern CA
<i>Mytilus edulis</i> (Blue mussel) (89)	0–27	5–33	differs between populations; some low-level throughout year; first in early summer, second in the fall	Atlantic coast, from Labrador to Cape Hatteras, NC
<i>Mytilus galloprovincialis</i> (Mediterranean mussel) (90)	8–25	10–33	Similar to <i>M. edulis</i> , but several weeks later when temperature is maximum	Mediterranean, Europe, Atlantic France and British Isles, Japan, East China to Korea, Australia, South Africa; southern CA to OR
<i>Mytilus trossulus</i> (Pacific blue mussel) (91)	0–29	4–33	July to September	Baltic Sea; west Coast, Central CA to AK; east Coast, Canadian Maritimes
<i>Ostrea lurida</i> (Olympia oyster) (92)	6–20	NA–25	Spring to fall: peaks in spring in south, mid summer in mid-range and north	Southeast AK to Baja California
<i>Protothaca staminea</i> (Littleneck clam) (93)	0–25	20–32	BC, Canada, January to March; AK, mid-July; southern CA, June	Aleutian Islands, AK to Cape San Lucas, Baja California
<i>Venerupis japonica</i> (Manila clam) (82)	13–21	24–31	Washington: once/year May–September; peaks in June/July	British Columbia to CA

the attached biomass with brushing or scraping, the bivalves should be transferred to clean, unfouled cages for the remainder of the exposure period.

6.11 Possible interferences associated with interpretation of tissue chemistry data include the use of inappropriate analytical procedures. It is critical to use the most appropriate method for each chemical analysis. Some commonly used methods, may not always be appropriate. For example, when measuring the suite of PAH-alkylated homologs, it is essential to use sufficient silica gel to clean up excess lipids in the sample. A more specific approach for these analyses developed as part of the Exxon Valdez oil spill assessment program included advanced methods specific to that group of researchers. These methods are recommended for bivalve tissues when source identification through chemical fingerprinting is necessary (94, 95, 96, 97).

6.12 Natural variability in the concentrations of chemicals of concern in water or sediment coupled with intermittent chemical discharges may increase the difficulty in interpreting exposure concentrations in these pathways. However, weekly measurements of chemicals in the water column coupled with measurements of bioaccumulation and growth have proven effective in explaining

the environmental significance of these variables (3, 18). In practice, it is usually difficult to sample with that frequency, and water samples are generally taken only at the beginning and end of the test. Since the variability in sediment chemistry is generally less extreme than in water, collecting sediment samples for chemical analysis at the beginning and end of test may be sufficient to characterize exposure conditions. However, sediments may also be highly variable on a small spatial scale (98).

6.13 In assessing effects of effluents with high organic loads, it is possible that the organic enrichment from the effluent will increase bivalve growth rates and make it more difficult to assess the adverse effects of associated chemicals. Differentiating between the positive effects of nutrient enrichment and the adverse effects of toxic chemicals is best accomplished by maximizing the number of stations in the assessment area, deploying caged bivalves at various depths, and maximizing the number of effects end points. The processes involved could be better characterized and understood by using various biomarkers in addition to the bioaccumulation and effects end points (99).

## 7. Hazards

7.1 Water and sediment might be contaminated with unknown concentrations of many potentially toxic materials. Any potentially contaminated water or sediment should be handled in a manner to minimize exposure of personnel to toxic compounds. Therefore, skin contact with all potentially toxic sediments and overlying water should be minimized by such means as wearing appropriate protective gloves, laboratory coats, aprons, and glasses particularly when washing equipment or placing hands into test water, effluents, sediment, or cleaning solutions. Respirators may also be necessary in some hazardous waste sites or during oil spills.

7.2 Water and sediment, particularly in effluent areas, might contain organisms that can be pathogenic to humans. Special precautions when working in these areas might include immunization prior to deployments and the use of bactericidal soaps after working in the water and touching sediments.

7.3 Use of ground fault systems is strongly recommended during measurements at the beginning and end of the tests where electronic equipment such as portable computers are used to record data electronically to help prevent electrical shocks because water is a good conductor of electricity.

## 8. Experimental Design

8.1 Field bioassays can be designed to provide either a qualitative reconnaissance or a quantitative assessment involving statistical comparisons of measured end points (that is, chemical concentration in tissues and effects end points) among stations. The object of a qualitative reconnaissance survey is to identify sites with the potential for bioaccumulation and associated biological effects. Qualitative surveys are often conducted in areas where little is known about contamination patterns. Quantitative assessments are conducted to test for statistically significant differences among stations.

8.2 Experimental design considerations, such as station location, number of stations per site, number of cages per station, and number of bivalves per cage, should be based on the purpose of the test and the procedure(s) used to analyze the results. Various experimental designs can be applied, with the most common used to:

(1) Compare bivalve tissue chemistry and growth at one or more stations to reference, background, or pre-test conditions.

(2) Compare bivalve tissue chemistry and growth among multiple stations to characterize patterns, trends, or gradients.

8.3 Experimental control of all test variables can be difficult to achieve in field tests that assess or monitor resident populations. The use of in-situ field bioassays allows the investigator to control the following: species; number; and size range of test animals, specific location(s) to be assessed, and exposure duration. Generally, the concentration of chemicals of concern and natural factors, such as temperature, salinity, dissolved oxygen, pH, current speed, and food supply, are not manipulated or controlled as they are in laboratory testing. However, ~~on a limited basis, chemicals and other additives~~ temperature could be pumped over the caged bivalves in much the same way as they are added to laboratory test tanks; chemicals could also be added as coated pellets or rods situated adjacent to the cages to administer controlled exposure. increased by heating dissolved oxygen by aeration, current speed by pumping, and food supply by adding nutrients. The intent of field bioassays is to determine chemical bioavailability and subsequent effects under natural, site-specific conditions, which includes intrinsic, site-specific variability. With an adequate number of stations, statistical testing can often identify the importance of these uncontrolled variables with respect to exposure and effects.

### 8.4 *Measurement End Points*

8.4.1 At a minimum, biological effects should be characterized by measuring survival and growth. Survival may not always be a very sensitive indicator of effects in bivalves (3), but it is an important parameter to monitor. Several factors can affect survival, including handling prior to test initiation and physical-chemical factors at the deployment stations. Survival can be easily quantified, although it is possible to have some individuals missing at the end of the test due to shell decomposition. Under some circumstances, more individuals may be present at the end of the test than at the start. This would most likely be due to the settlement of juvenile bivalves during the course of the test. This can easily be accounted for as new recruits should be smaller than the test bivalves. All recruits should be removed prior to determining survival and assessing effects end points. Only effects measurements from surviving bivalves should be used to calculate summary statistics. It is possible for shells to stick together due to mucilaginous material or sediment within the shells, prohibiting a precise determination of death. Thus, all dead bivalves may not be identified until the tissue removal process when the shells are opened to reveal the internal tissues.

8.4.2 Growth is a sensitive sublethal effects end point that is easy to measure and is recommended for all field bioassays. It is

generally more sensitive than mortality, and reductions in growth have been related to adverse effects on bivalve populations (1). As many growth end points as are practical should be measured for assessing growth in a weight-of-evidence approach. For example, it has been shown that shell growth and tissue growth are decoupled. Measuring only one of these end points could provide misleading results and lead to a spurious interpretation of environmental effects on growth (53,54). Growth end points include, but are not limited to, whole-animal wet weight, shell length, tissue weight, shell weight (18). Whole-animal wet-weights and shell lengths are nondestructive measurements and can be made multiple times over the course of the exposure period. At a minimum, whole-animal wet-weights and shell lengths should be measured at the beginning and end of the test. Since tissue weights and shell weights provide a different perspective on animal health and may be related to different stressors, they should also be measured at the beginning and end of the test (18). Because these measurements are destructive, beginning-of-test tissue and shell weights for bivalves to be deployed can be estimated from a subsample of the bivalves distributed to the individual cages. The bivalves used to estimate beginning-of-test tissue and shell weights should be within the same size range as those identified for deployment. Because the initial tissue and shell weights are based on a subsample of the test population, the change in these metrics over the test period is an indirect determination and has some uncertainty. However, tissue and shell weights can provide the most discriminating measurements under certain conditions, particularly when growth rates are low (18).

8.4.3 Although tissue dry weights are less variable than wet weights, drying the tissues has some limitations. (1) It is more time consuming to dry all the tissues and make the weight measurements. (2) In a combined bioaccumulation and biological effects test, the same wet tissues can be used for chemical analysis and wet-weight measurements (drying tissues may destroy organic chemicals, and limit their ability to predict bioaccumulation potential). (3) A wet-weight approach has been used successfully (18), and may provide better correlations with other growth metrics. Nevertheless, if additional testing clearly demonstrates an advantage to measuring dry weights, or if particular studies require more emphasis on the accuracy of tissue weight measurements, it would be relatively simple to alter the procedures accordingly.

8.5 *Reference Stations*—The use of one or more reference stations may be used for field bioassays with caged bivalves. It is the responsibility of the investigator to determine the need for reference station(s) based on the experimental design. It may be difficult or problematic to identify a true reference in the field because of the variability in field conditions and the influence of natural factors on site-specific conditions. If reference stations are used, the physical and natural factors (that is, temperature, salinity, dissolved oxygen, vegetation, and currents) at the reference station(s) should be as similar as possible to those conditions at the area under investigation. Multiple reference stations may help account for natural differences and variability among uncontaminated areas. It may be more useful to employ a gradient design with decreasing chemical gradients in bivalve tissue chemistry associated with changes in growth rate rather than comparing treatments to reference conditions or upstream versus downstream sites.

8.56 Natural population of bivalves could also be used for comparative purposes, but these comparisons should be made cautiously because there is evidence that caged bivalves can have different growth rates and different rates of accumulation than natural populations under certain conditions (3, 11, 12). It would be useful if growth rates of natural populations and caged bivalves were compared, if practical.

8.67 *Statistical Design*—Field bioassays with caged bivalves can be used to support a variety of statistical designs. The experimental design is a function of the technical and environmental issues to be answered as well as the most appropriate statistical design for analyzing the data. The level of replication is a function of desired power and confidence. The following null hypotheses can be used to determine statistical differences in bivalve bioaccumulation and associated biological effects among stations as well as relationships between tissue chemistry, sediment or water chemistry, if measured, and measured effects:

(1) *Null Hypothesis #1*—There is no difference in bioaccumulation of chemicals of concern (as determined by tissue burdens) between test and reference station(s),

(2) *Null Hypothesis #2*—There is no difference in effects between test and reference station(s),

(3) *Null Hypothesis #3*—There is no relationship between effects end points in bivalves and tissues, water, or sediments containing chemicals of concern among stations.

(4) *Null Hypothesis #4*—There is no relationship between bioaccumulation and associated biological effects with distance from the suspected chemical source.

The preceding null hypotheses can also be used when it is appropriate to pool the stations to allow comparisons among sites. It may also be appropriate to apply these hypotheses to gradient designs in both horizontal and vertical planes in the water column or in bottom sediments.

8.78 *Replication*—The individual bivalves or the cage may be considered as the experimental unit. It is the investigator's responsibility to define the experimental unit and level of replication, which are appropriate for the study design. Additional guidance on statistical approaches can be found in other ASTM standards (E 1847 and E 1191). The distance between stations, or cages, is a function of the size of the area under investigation, the expected gradient or change in monitoring parameter(s), and the expected variability in conditions. Typically, stations can be placed 50 to 500 m apart. However, stations can be closer together, or further apart, as determined during development of the study design and hypotheses.

8.78.1 For the exposure assessment, a chemical replicate may be formed by combining the tissues of all living bivalves from one cage (see 9.3). Compositing may be necessary because, in most cases, individual bivalves do not contain sufficient tissue for chemical analysis. The cage can be used as a way to identify the bivalves to be combined for a chemical replicate. The number

of chemical replicates prepared for each station depends on the level of replication desired for the bioaccumulation assessment. If statistical comparisons are desired, a minimum of three replicate tissue samples for each station is recommended. The number of bivalves required for each tissue sample is a function of the tissue mass requirements for the chemical analyses being performed and the tissue mass of the individual bivalves. The analytical laboratory performing the chemical analyses should be contacted to identify the amount of tissue required for each analysis. For example, if the analytical laboratory requires a minimum of 50 g wet tissue, and the average individual tissue weight is 0.5 g wet, then a minimum of 100 bivalves will be required for each chemical composite (that is, 100 bivalves per cage). With larger bivalves there may be sufficient tissue to conduct chemical analyses on individuals, particularly if only a few chemicals are being analyzed. This approach could improve the discriminating power of the assessment.

**8.78.2** For the effects assessment, each individual bivalve may be considered a replicate, although concerns about individuals in the same experimental unit have been noted (Guide E 1847). The bivalves within a predetermined size range are assigned to cages (see Section 11), and the cages are randomly assigned to stations. Independence among bivalves within each cage is assumed. In addition to the tissue chemistry biomass requirements, the minimum number of bivalves per cage should also consider the following with respect to effects endpoints: (1) the expected variance within cages, (2) the expected variance between cages, and (3) either the maximum acceptable width of the confidence interval on a point estimate or the minimum difference that is desired to be detectable using hypothesis testing.

**8.89** *Statistical Analyses and Data Interpretation*—The calculating procedure(s) and interpretation of results should be appropriate to the experimental design. Procedures used to calculate results of these field bioassays can be divided into two general categories: those that test hypotheses regarding differences among stations, and those that establish relationships along suspected chemical gradients or between bioaccumulation and growth in the test organisms. No procedure should be used without careful consideration of (1) the advantages and disadvantages of various alternative procedures, and (2) appropriate preliminary tests such as those for outliers and heterogeneity. Preprocessing of data might be required to meet the assumptions of the analyses. All parameters measured at the end of the test (that is, whole-animal wet-weight, shell length, tissue weight, shell weight, and chemical concentrations in tissues) can be statistically analyzed. Summary statistics (for example, mean and standard deviation) can be calculated for each of these parameters on a station-by-station basis. The appropriate statistical test is a function of experimental design, hypotheses, and measurement end points. It is the investigator's responsibility to identify the appropriate statistical tests. In general, ANOVA and multiple comparison tests are used for hypothesis testing and comparison among stations. Linear regression analysis is generally used to establish relationships between bioaccumulation and growth end points along suspected chemical gradients and to establish relationships between bioaccumulation and growth. If statistical differences are found, a multiple range test can be used to determine which stations are different from the others. A textbook on statistical analyses of biological data can be referenced for appropriate tests and procedures (**100, 101, 102**).

**8.89.1** Power analyses performed on data from caged bivalve studies in Alaska indicate that between 100 and 300 mussels per station are sufficient to detect differences in weight on the order of 0.2 and 0.1 g wet, respectively. An environmental significance, or likely adverse effect to the community, is expected when both a statistically significant difference is observed ( $\alpha = 0.05$ ) and there is a 10 to 25 % absolute difference between the test and reference/control station(s) (**5**).

**8.910** *Test Duration*—For most studies, bivalves should be exposed to site-specific conditions for a minimum of 30 days. An exposure period of less than 30 days is not generally recommended unless the chemicals of concern are low molecular weight organic compounds, such as some PAHs. Equilibrium for most other chemicals, such as metals and high molecular weight organic compounds, is generally achieved in marine and freshwater bivalves within a period of approximately 60 to 90 days (**3, 11, 12, 15, 66, 103, 104, 105, 106, 107, 108**). If both exposure and effects end points are being measured, it may be advantageous to continue the test for 60 to 90 days to facilitate chemical equilibrium and provide sufficient time to allow adverse effects to manifest themselves. Extending the exposure period may also increase the ability to detect statistically significant differences among effects end points. Although unlikely, it is possible that deployment of caged bivalves in or on bottom sediments may reduce concentrations of some chemicals. This may be particularly important in very small areas with restricted circulation where bivalves are removing chemicals from sediment. Consistent sediment conditions during the deployment period can be verified by sampling the sediment before and after deployment. It is the responsibility of the investigator to verify concentrations of chemicals in sediment before, during, and after deployment if this is an issue of concern.

## 9. Apparatus

**9.1** *Facilities*—Sources of water and power and the ability to be protected from rain, snow, and wind can be of considerable help in sorting the animals at the beginning of the test, making the appropriate measurements, and removing tissues for chemical analysis at the end of the test. Preparations can be made outdoors, but inclement weather can interfere with making accurate measurements. The portable analytical balance is particularly sensitive to wind although some protection can be provided by a wind barrier around the entire area, such as a lean-to, or a smaller barrier such as a box to protect the balance. Making weight measurements aboard boats or floating piers is not recommended, unless the measuring devices are specifically designed for use on unstable platforms. Length measurements made with calipers are not affected by the instability associated with boats or floating piers.

**9.2** *Construction Materials*—Equipment such as cages, predator mesh, holding tubs, and facilities ice chests, that contact the test water, sediment, and organisms should not contain substances that can be leached or dissolved by aqueous solutions in amounts

that can adversely affect test organisms or be accumulated in their tissues. In addition, equipment that contacts test water, sediment, and organisms should be chosen to minimize sorption of test materials from water. Glass, Type 316 stainless steel, nylon, high-density polyethylene, polycarbonate, and fluorocarbon plastics should be used whenever possible to minimize dissolution, leaching, and sorption, except that stainless steel should not be used in saltwater. Concrete may be used for cage anchors and rigid plastics (that is, PVC) may be used for weights cage frames. Plastic Frames and for frames, but they mesh bags should be soaked before use, preferably in flowing fresh or seawater, for at least 24 h to remove water soluble and volatile chemicals. Mesh bags, tubes, or trays used to create the compartmentalized cages for holding the mussels during deployment should be made from high-density polyethylene, polycarbonate, or fluorocarbon plastic. Plastic cable ties have many applications during cage construction, such as separating the individual bivalves when mesh bags are used and attaching cages to deployment moorings and lines. Plastic cable ties should not contain metal stops as these can corrode and break upon exposure to water. This corrosion can result in detachment or addition of chemicals. Brass, copper, lead, cast iron pipe, galvanized metal, and natural rubber should not contact water, sediment, or test organisms before or during the test.

### 9.3 Cages:


~~9.3.1 Cages with individual compartments are recommended for field studies with caged bivalves. The~~

9.3.1 The basic concept behind the cage design is to maximize mesh size to maximize water flow to the test animals yet maintain animals. This is accomplished by using a mesh size large enough to maximize flow but small enough to contain the test animals. Cages with individual compartments are recommended for field studies with caged bivalves. The separation of individuals into individual compartments allows equal exposure to each bivalve (Fig. 1C). Compartmentalization facilitates tracking individuals throughout the test and eliminates the need to mark or notch individuals. Compartmentalization permits multiple growth measurements on individuals, ensures that an accurate record of measured end points can be maintained on individuals, and facilitates conducting tissue chemistry analysis on individuals if the individual bivalves contain sufficient biomass. Recording measurement data on an individual-by-individual basis increases the statistical power of the test. Each of the measurement end points, including tissue chemistry, can be paired during statistical analyses.


9.3.2 In its simplest form, in-situ field tests can be conducted with bivalves held in cages without compartments as shown in Fig. 1A (109). This approach is not recommended because it limits the ability to make multiple measurements on the same individuals throughout the course of the test. There are techniques for numbering individuals (110, 111), but this may be prohibitively time consuming if large numbers of animals are being caged. Numbering with different glues and epoxies could also introduce other potentially toxic chemicals. Cages can also be rigid with fixed compartments (Fig. 1B), as in plastic trays and wire baskets. Rigid cages with fixed compartments have been used in freshwater (112, 113) and marine (3) environments. Cages can also be a combination of flexible mesh material with compartments attached to a rigid frame (Fig. 1C), as with mesh bags attached to a PVC frame. This approach has been used in freshwater (3), estuarine (68), and marine habitats (15). The flexible mesh bags can also be attached to heavy plastic mesh that serves as protective cage and an attachment point for the bags. The mesh bags used to hold the bivalves are created from tubular oyster culch netting similar to that used in bivalve aquaculture. The bivalves are separated within the mesh bags by placing a plastic cable tie or other restricting device between individuals. Different cage designs have also been tested to compare with the performance of natural bivalve populations. These include rigid cages with and without compartments, corrals that limit the movement of sediment-dwelling bivalves, and leashes where monofilament lines were glued to each bivalve shell (114).

9.3.3 The final dimensions of the deployment cages depend on the size of the individual test organisms and the number of organisms per cage. One advantage of using the flexible mesh bags and a PVC frame is that the size of the individual compartments and the overall cage size can be easily adjusted. Sufficient space should be provided in each compartment to allow test animals to open their valves and grow during the exposure period; the amount of space depends on the species used, the size of individuals at the start of the test, and expected increases in growth over the deployment period. For rigid cages, investigators should make the individual compartments large enough to accommodate expected growth during the test. A 6-in. (approximately 15-cm) diameter mesh material is recommended for smaller smoothed-shelled species like mussels and clams because there is less excess mesh at the point of constriction. For larger bivalves with rough shells and irregular shapes, such as oysters, it may be necessary to use a tubing of larger diameter. Because the flexible mesh is tubular in form, it is not necessary to adjust the width/height dimensions. The length of each compartment in the mesh bag (that is, the distance between constricting cable ties) should be large enough to accommodate valve opening and expected growth during the test. The mesh bag should be long enough to accommodate the desired number of bivalves per bag plus sufficient material to allow secure attachment to the PVC frame. Approximately 30 cm of mesh netting on either end of the bag is generally sufficient for attachment to a PVC frame constructed from 3/4-in. (approximately 1.90-cm) material. The PVC frame should be approximately 5 cm longer than the space occupied by the bivalves positioned in the mesh bag. The width of the frame should be about 5 cm greater than the distance occupied by all mesh bags to be attached to the frame when laid side-by-side.


9.3.4 If PVC cages are to be deployed on top of sediments, pushed a short distance into the sediments, or positioned where neutral buoyancy is desired, the PVC pipe should be drilled approximately every 24 cm with a 1/4-in. (approximately 0.64-mm) hole to allow water to enter the pipe and remove trapped air. The corners of the frame should not be ~~drilled to eliminate drilled.~~ Drilling the potential for weakening corners could weaken the overall structure of the frame. For water column deployments,



A. Mesh basket (front view)



B. Compartmentalized mesh tray (top view)



C. PVC frame supporting mesh bags (side view)

FIG. 1 Possible Cage Types for In-situ Field Tests With Caged Bivalves

flotation can either be increased or decreased depending on whether the PVC frames are drilled to allow a water ballast or left undrilled to add extra flotation.

9.4 *Cage Deployment Configuration*— The methods used to deploy cages and the type of mooring system depends on the experimental design identified for the specific media being assessed and substrates of opportunity. It may be useful to conduct a reconnaissance of the deployment area prior to setting out the cages to allow identification of potential deployment impediments and potential interference from the public. If floating or fixed piers are available in the assessment area, they could provide a potentially effective substrate for attaching bivalve cages. Figs. 2-5 provide various deployment configurations, and for simplicity, only rigid cages are shown. The PVC frames supporting bivalves in mesh bags can also be used in the same deployment schemes. Fig. 2 shows caged bivalves attached to floating (3) and fixed piers (5). Under most circumstances structures such as piers may not be available and open-water, nonstructural deployments should be used as shown in Fig. 3A (3) and Fig. 3B. A more direct assessment of bottom sediment is possible with fixed bottom deployments as shown in Fig. 4A (3, 67) and Fig. 4B (5, 115)). Caged bivalves can be placed directly on bottom sediment or on legs used to raise the cages above the sediments. Cages with legs can also be used to stabilize the unit and maintain position in high energy areas such as the intertidal zone. The most sophisticated assessments include a gradient design with cages placed at multiple depths and distances from suspected sources, as shown in Fig.

## Pier Suspension




FIG. 2 Pier Deployments: Floating or Fixed

## Single Depth in Water Column




FIG. 3 Open Water, Nonstructural Deployments

## Fixed Bottom Placement




FIG. 4 Fixed Bottom Deployments

5. Each of the preceding deployment configurations uses rigid or flexible compartmentalized cages. Similar deployment configurations have been used with rigid and flexible un-compartmentalized cages (21, 61, 116, 117, 118). Placing cages along suspected chemical gradients in three dimensional space helps to identify not only potential sources of chemicals, but the relative contribution of chemicals in the water column and sediments based on chemical concentrations measured in bivalve tissues.

9.4.1 *Water Column Assessment*—To evaluate the bioavailability and potential effects of chemicals within the water column, cages can be suspended from a fixed mooring (that is, floating pier, piling, or other fixed structure), suspended within the water column by attachment to a line that has an anchor or weight on one end and a surface or subsurface buoy attached to the other end, or fixed in the water column by attaching legs to the cages and pushing the legs into the sediments to hold the cages in place. Depending on the species of bivalve used, bivalves in cages deployed directly on top of the sediments can be used to assess chemicals within the water column as well. Factors that should be considered during the deployment of cages for surface water

## Gradient Design: Multiple Depths and Distances




FIG. 5 Compartmentalized Cages Attached to Line at Three Depths (With Surface Float and Bottom Anchor) and at Multiple Distances from a Suspected Containment Source

assessments include change in tidal height (that is, to ensure the cages are at the desired depth during both low and high tides), ~~slope of bottom material~~ slope (that is, to ensure the cages do not slide down a steep slope during the exposure period), and boating activity and recreational activity in the vicinity of the cages (that is, to avoid cages being removed by or tangled within propellers). Floats should be appropriate to accommodate the weight of the line plus cages at the depth of deployment. This type of water column monitoring has been used to evaluate PAHs, dioxins and methylmercury, tributyltin (4, 18) and pulp mill effluents in marine environments. Similar methods have been used to assess freshwater sites (114).

9.4.2 *Sediment Assessment*—One way to evaluate chemicals associated with surficial sediments is to position the cages directly on top of the sediments. This facilitates positioning the test species in the upper layers of sediment where most benthic organisms are commonly found and from which sediments are most commonly collected for chemical analysis (Guides E 1391 and E 1525) as well as for use in laboratory bioaccumulation tests (Guide E 1688) and laboratory toxicity tests (Guide E 1367 and Test Methods E 1706). Allowing the bivalves to bury themselves is generally less stressful than forced burial, and allows the cages to be positioned without divers in water deeper than 2 m. Placing cages directly on top of the sediments and allowing the bivalves to reposition themselves has been used successfully in freshwater (3, 67) and marine (68) environments. The bivalves gain exposure to sediments as the sediments infiltrate the mesh material. Although confined within the mesh material, bivalves can work themselves into the sediment if sufficient room within their compartments is provided. In some cases it may be necessary to push the cage into the sediment to ensure exposure to chemicals associated with those sediments. This approach has been used before (119), although without compartmentalized cages. However, forcing the cages and bivalves into the sediments may induce stress, resulting in high mortalities. Forcing the cages into the sediments, or digging out sediments to bury the cages can disturb the integrity of the sediments and alter natural biogeochemical processes. Anchors, rebar, or cages with legs can be used to ensure the cages remain at the desired position. Anchors or cement blocks can be attached to the sides of the cage with line. Rebar can be used as a weight and strapped to the side of the cage or it can be bent into a “U,” and pushed over the cage into the sediment to secure the cage in place. However, rebar should be coated with rubberized coatings or covered by plastic bags to prevent potential metal exposure. The cage can be constructed with legs (that is, like a table) and the legs pushed into the sediment to secure the cage in position. Factors that should be considered when deploying cages on top of the sediments include presence of natural vegetation, type of substrate, boating traffic, and recreational activity.

## 10. Test Organisms

10.1 *Species*—Many bivalve species have been used for assessing chemical bioavailability or effects in marine, estuarine, and freshwater environments (Table 3). The most widely used approach has been monitoring tissue chemistry of resident populations. The existing information on species life history, ability to accumulate chemicals of concern, and physiological effects can be applied to in-situ field bioassays. Species selection should be made carefully, considering conditions at the natural habitat of the species and natural factors (that is, temperature, salinity, dissolved oxygen, and pH; Table 2) at the area under investigation, presence in the area under evaluation, documented ability to accumulate chemical(s) of concern, documented sensitivity to chemical(s) of concern, life history (that is, spawning cycle, life-stage requirements, and threatened or endangered status), availability, and ease of handling in the field. Ideally, species or genera with wide geographic distributions should be selected, so that test results can be compared among different sites and different test conditions. Depending on the question being asked, it may be most important to select species that are found, have been found, or could be found in the assessment area. Species used should be identified with an appropriate taxonomic key, and identifications should be verified by a taxonomic authority wherever possible. It may be necessary to conduct a pilot study to determine if the test animals can survive under the environmental conditions at a particular site. The U.S. Fish and Wildlife Service (120) has identified the following mussels as possible surrogate species for in-situ field testing instead of threatened and endangered species in the Virginia area:

- (1) Rainbow mussel (*Villosa iris*),
- (2) Snuffbox mussel (*Epioblasma triquetra*),
- (3) Pimpleback mussel (*Quadrula pustulosa*), and

**TABLE 3 Partial List of Bivalves Used in Transplant Studies for Monitoring Marine and Freshwater Environments**

NOTE—Many species can be used in estuarine environments due to their ability to tolerate a wide range of salinities. “\*” indicates species used in studies with compartmentalized cages; most other species have been deployed in non-compartmentalized cages, although a few benthic studies did not employ cages of any kind.

Marine Species			
Mussels	Oysters	Clams	Scallops
<i>Arca zebra</i> (121)	<i>Crassostrea gigas</i> (130)	<i>Anadara granosa</i> (138)	<i>Aequipecten opercularis</i> (147)
<i>Modiolus</i> (122)	<i>Crassostrea angulata</i> (131)	<i>Cerastoderma edule</i> (139)	<i>Amusium pleuronectes</i> (148)
<i>Mytilopsis sallei</i> (123)	<i>Crassostrea virginica</i> * (132)	<i>Chione stutchburyi</i> (140)	<i>Argopecten irradians</i> (149)
<i>Mytilus edulis</i> * (124)	<i>Crassostrea rhizophorae</i> (133)	<i>Macoma nasuta</i> (141)	<i>Argopecten purpuratus</i> (150)
<i>Mytilus galloprovincialis</i> *(125)	<i>Ostrea angasi</i> (134)	<i>Macoma balthica</i> (119)	<i>Chlamys varia</i> (151)
<i>Mytilus trossulus</i> * (126)	<i>Ostrea edulis</i> (135)	<i>Macoma inquinata</i> (142)	<i>Crassodoma gigantea</i> (152)
<i>Mytilus californianus</i> *(127)	<i>Ostrea lurida</i> (136)	<i>Mercenaria mercenaria</i> (143)	<i>Lima hians</i> (153)
<i>Perna viridis</i> * (128)	<i>Saccostrea commercialis</i> (137)	<i>Mya arenaria</i> (144)	<i>Pecten maximus</i> (154)
<i>Perna bicolor</i> (129)		<i>Scrobicularia plana</i> (145)	<i>Placopecten magellanicus</i> (155)
		<i>Spisula solidissima</i> (146)	
		<i>Venerupis japonica</i> (141)	
		<i>Venerupis staminea</i> (141)	
Freshwater Species			
Mussels	Clams/Cockles		
<i>Actinonaias ligamentina</i> (156)	<i>Anadara tpeziium</i> (177)		
<i>Actinonaias pectorosa</i> (156)	<i>Corbicula fluminea</i> * (178)		
<i>Amblema plicata</i> (157)	<i>Corbicula japonica</i> (179)		
<i>Amblema perplicata</i> (158)	<i>Corbicula manilensis</i> (180)		
<i>Anodonta anatina</i> (159)	<i>Musculium transversum</i> (181)		
<i>Anodonta cygnea</i> (160)	<i>Rangia cuneata</i> (182)		
<i>Anodonta grandis</i> (161)	<i>Sphaerium striatinum</i> (183)		
<i>Anodonta implicata</i> (162)	<i>Sphaerium simile</i> * (184)		
<i>Anodonta piscinalis</i> (70)			
<i>Anodontites trapesialis</i> (163)			
<i>Dreissena polymorpha</i> *(164)			
<i>Elliptio complanata</i> * (165)			
<i>Epioblasma torulosa rangiana</i> (166)			
<i>Epioblasma triquetra</i> (166)			
<i>Fusconia subrotunda</i> (156)			
<i>Hydridella menziesi</i> (167)			
<i>Lampsilis higgins</i> (168)			
<i>Lampsilis radiata</i> (169)			
<i>Lampsilis ventricosa</i> (170)			
<i>Lemiox rimuosus</i> (171)			
<i>Margaritifera falcata</i> (172)			
<i>Medionidus conradicus</i> (156)			
<i>Proptera alata</i> (173)			
<i>Proptera capax</i> (174)			
<i>Pyganodon grandis</i> (42, 43, 44)			
<i>Quadrula quadrula</i> (175)			
<i>Unio pictorum</i> (160)			
<i>Villosa nebulosa</i> (156)			
<i>Villosa vanuxemensis</i> (156)			
<i>Westralunio carteri</i> (176)			

(4) Tennessee pigtoe mussel (*Fusconaia barnesiana*). A special caution is required to prevent the introduction of exotic species (for example, zebra mussels (*Dreissena polymorpha*) and Asiatic clams (*Corbicula fluminea*) in freshwaters and clams (*Potamocorbula*) in marine waters). To prevent the spread of unwanted species, it must be verified that they already exist in the area of concern before using them in an in-situ field bioassay. Zebra mussels, Asiatic clams, and Potamocorbula are good candidates for in-situ field bioassays and have been used successfully (for example, (49, 50, 51, 52), but these species should be used with extreme caution and consultation with appropriate regulatory agencies. The introduction of zebra mussels can be particularly harmful because of their potential effects on freshwater mussels and surrounding environments. Other unwanted introductions include parasites or diseases (*Perkinsus marinus*) from infected bivalves such as oysters. Similarly, unwanted pests could be introduced from microscopic attached forms on the shells of transplanted bivalves such as the freshwater weed *Hydrilla hydrilla*. A potential problem that is unique to freshwater bivalves is the introduction of glochidia stages that are parasitic on fish for a portion of their life cycle. Although this is common in areas where the freshwater bivalves naturally reside, potential effects should be considered during the planning of in-situ field bioassays (for example, see 10.2.1).

10.2 *Commonly Used Taxa*—The environmental requirements (Table 2) and sensitivity of new bivalve test species should be established before they are widely used in field tests. The sensitivity and bioaccumulation potential of a prospective new test species could be compared with a more commonly used species to establish its relative utility. This can be accomplished in side-by-side transplants. Monitoring variations in water quality parameters and sediment characteristics (that is, particle size, organic enrichment, and sulfides) may help distinguish the effects of these parameters from the effects of chemical exposure. The taxa most commonly used in in-situ field bioassays are described as follows:

10.2.1 *Mytilus* sp. is an intertidal bivalve that has been successfully used in transplant studies since the late 1970's (46). The sensitivity of this species to salinities less than 10 ppt limits its use to testing marine and estuarine areas. However, the large data base that has been developed for the response of *Mytilus* sp. to a variety of habitats and chemicals establishes its usefulness as a test species as well as a reference species for comparing the sensitivity of other species. Species of the genus *Mytilus* are widely distributed on both coasts of North America (90).

10.2.2 *Corbicula fluminea* is a freshwater clam that has been used extensively in field transplants and laboratory studies (112, 113, 185, 186, 187, 188). Numerous laboratory studies have also been conducted on this species and several symposia have been conducted on their biology and ecology.

10.2.3 *Elliptio complanata* and *Pyganodon grandis* (formerly *Anodonta grandis*) are freshwater unionid mussels that have been used extensively for monitoring water column and sediment exposures in northern parts of the United States and in Canada (3, 42, 43, 44, 189). Numerous laboratory studies have also been conducted on these species.

10.2.4 *Dreissena polymorpha*, a freshwater mussel, is a relative newcomer to bivalve field bioassays. However, the rapid proliferation of this nuisance species has increased the number of laboratory and field studies that have been conducted on bioaccumulation and growth (49, 50, 51, 52). It is particularly important to limit unwanted introductions of nuisance species into areas where they are not already found.

10.2.5 *Macoma* sp. the only commonly used facultative deposit feeder, is a marine or estuarine clam that has been used extensively in laboratory and field studies to assess bioaccumulation and growth (11, 12). It is commonly found in many environments on several coasts. It has been successfully transplanted in many different areas, and there are a number of supporting laboratory studies.

10.2.6 *Rangia cuneata* is an estuarine clam that can tolerate freshwater conditions (182, 190, 191, 192, 193). It has been used in a number of field transplants as well as laboratory studies.

10.2.7 *Crassostrea* sp. is an oyster that has been used extensively in transplant studies in marine and estuarine studies (194, 195). Oysters survive and grow better than marine mussels at lower salinities and accumulate many chemicals such as tributyltin and copper by about a factor of two above mussels. The shells of *Crassostrea* and other oyster species are usually more difficult to measure because of the irregular shell shape, sharp edges and protrusions.

10.3 *Size and Age of Test Organisms*— All bivalves used in an in-situ field study should be from the same age class and as uniform as possible in size. Age class is more difficult to determine when obtaining specimens from the wild as opposed to culturing facilities because wild populations are a composite of several age classes, with different individuals growing at different rates. In most bivalve species, size is a function of age, so if individuals from the same age cohort are selected, they will be within a fairly uniform size range. Under adverse conditions (that is, crowding, exposure to chemicals, or exposure to unfavorable natural factors), some individuals may grow very little. The age or size class of the prospective species should be chosen so that sensitivity to chemicals or bioaccumulation potential is not affected by state of maturity or reproduction. It is recommended that specimens in a sub-adult age class be used because this age class has the greatest potential for growth of somatic tissue, reproductive tissue, and shell. If adult specimens are used, the study should not be conducted during active spawning. This prevents loss of accumulated lipophilic chemicals. Investigators should note the reproductive state of the test animals including degree of gonad development by observation, mass, or volume measurements. They should also note whether spawning occurred during the beginning-of-test or end-of-test measurements.

10.3.1 Bivalves can be sorted according to size and counted to determine whether sufficient numbers have been collected in the desired uniform size range. Shell length or whole-animal wet-weight should be used to select individuals for use in a field bioassay.

Although whole-animal wet-weights provide a more accurate measurement of animal size, shell lengths provide a rapid and quantitative sorting method. Shell length should be determined with vernier calipers with a measurement accuracy of 0.1 mm. Whole-animal wet-weights should be determined with an analytical balance with a measurement accuracy of 0.01 g. The final size range used should be based on the maximum number of animals in the minimum size range. As a starting point, it is recommended that the size range used in the test be approximately 5 to 10 % of the average maximum size of the species (that is, for *Mytilus* a 5 to 7-mm size range is suggested as the maximum shell length this genus is about 70 mm). By minimizing the size range of individuals, the variability in bioaccumulation and associated biological effects will also be minimized. The absolute size range used for a given species will depend on the size of the species and the availability of specimens. The decision to use juveniles, sub-adults, or adults depends on the experimental design. There is a tendency among many bivalve species for the smallest animals to grow at the greatest rates and accumulate the highest concentrations of chemicals. Various bivalve life stages have been proposed as part of an integrated environmental assessment (196). There are ASTM standard guides for a saltwater bivalve embryo test (Guide E 724), bioconcentration tests with adult saltwater bivalves (Guide E 1022), and bioaccumulation of sediment-associated chemicals (Guide E 1688). A freshwater glochidia test is also being developed (197). While bivalve larvae are often assumed to be the most sensitive life stage, there is a growing body of evidence suggesting they are not necessarily more sensitive than adults. Adults may be more sensitive than juveniles under certain conditions (2, 3, 11, 198, 199).

10.4 *Source*—Bivalves can be obtained from either natural populations or from culturing facilities. All individuals used in a field study should be from the same population, because different populations of the same species might have different sensitivities to or bioaccumulation capacities of the same chemical. Bivalves can be collected from wild populations in an uncontaminated (that is, chemicals present at concentrations lower than at the area under investigation) area although it may be easier to purchase species from field grow-out or laboratory culture facilities. The advantages of using cultured or farmed animals is that the genetic and environmental history of the test animals is well known, and the assurances of being uncontaminated are greater. A sample of the prospective test bivalves should be measured for contamination in their tissues, particularly for the chemicals of concern. Collecting permits for field collected bivalves might be required by some local and state agencies.

10.5 *Number of Specimens*—The number of bivalves collected should account for dead or dying individuals, individuals injured during handling, and the ability to minimize the size range as much as practical. Therefore, the number collected should be equal to the total number required for deployment at each of the stations plus the number required for the baseline (that is, beginning of test) tissue chemistry measurements, plus approximately 20 to 50 % more individuals to account for mortalities and size variability. Between 100 and 300 animals may be sampled at the beginning of the test to provide the initial tissue and shell weight estimates. These same individuals can be used for baseline chemical analysis. If composite tissue samples are required, see 8.78.1 for methods to determine number of animals per composite. It may be convenient to use a sample size equal to the number of individuals in each cage. Additional guidance on statistical approaches can be found in ASTM Practice E 1847 and Guide E 1191.

10.6 *Collection*—Natural populations should be collected with methods appropriate to their distribution. For intertidal marine species or freshwater species in shallow water, populations can be collected by hand. For subtidal marine species or freshwater species in deeper water, SCUBA or a small biological dredge can be used for collection. Infaunal bivalves can be separated from sediment by gentle sieving. Sieves and containers used to collect and transport bivalves should be marked “live only” and should never be used for working with formalin or any other toxic materials. Water used for sieving should be at the same temperature and salinity as bottom water at the collection site. Some species of marine and freshwater mussels produce byssal threads as an attachment mechanism. Particular care should be used when removing mussels from substrates to which they have attached to avoid damage to the byssal glands, an internal organ that secretes the byssal threads. Damage or removal of the byssal gland can lead to mortality. A knife or scissors should be used to remove the mass of byssal threads visible on the outside of the mussel shell. This process will reduce the possibility of injury to the mussel. All epiphytic growth should be removed from the exterior of the bivalve shells. Plant or animal growth can usually be removed by hand; a soft brush or scraper may be required to remove barnacles, tube worms, or other tenacious organisms.

10.7 *Handling*—Test organisms should be handled as little as possible. When handling is necessary, it should be done carefully, gently, and quickly so that specimens are not unnecessarily stressed. Every effort should be made to maintain bivalves in well aerated, flowing water for as long as possible between collection, sorting, and deployment procedures. The water used during the holding period(s) should be from a known source and free of chemical contamination. When transporting bivalves over great distances that require extended periods of time, it is better to keep them moist and cool than to maintain them in water that could become stagnant and low in dissolved oxygen. Bivalves can be kept moist and cool by placing the specimens in an ice chest with either wet ice or frozen gel packs on the bottom. Newspaper, paper toweling, or cloth toweling should be placed between the specimens and ice to prevent direct contact. Wet towels can also be placed over the specimens to provide additional moisture.

10.8 *Holding*—Test organisms should be deployed as soon as possible and holding times minimized. If necessary, test organisms can be acclimated to water quality conditions at the deployment site, as identified in 6.5. In larger studies where it may be difficult to collect a sufficient number of specimens in one day, a laboratory or field site relatively free of contamination can be used as a holding facility while the remainder are collected.

10.8.1 Some infaunal bivalves may require holding in sediment until initiating the test. Supplementary feeding for laboratory held specimens should not be necessary if holding is less than one week; specimens held in the field will continue to feed on natural sources of food.

10.9 *Animal Quality*—All bivalves used in a test need to be of acceptable quality. Before initiating a test, a qualified bivalve taxonomist should be consulted to ensure that the animals collected are all of the same species. This is particularly important with some freshwater bivalves where species differences may be extremely difficult to determine based on shell morphology. Even in the genus *Mytilus* there are subtle differences that may not be obvious, particularly in areas where two species could be found together (200, 201, 202, 203, 204) .

10.10 Although it is extremely difficult to identify healthy animals when the shell is closed, gaping animals that close very slowly or do not close at all should not be used. A putrefied smell emanating from the batch of test bivalves indicates one or more dead specimens. Dead specimens should be removed.

## 11. Field Procedures

11.1 *Test Initiation: Presort*—The first step is to sort all bivalves into size groups, with each size group in its own container. As indicated in 10.2, sorting can be based on either shell length or whole-animal wet-weight. Sorting allows identification and numeration of bivalves in a uniform size range. All bivalves originally collected should be initially retained until the test is initiated to ensure a sufficient number of individuals. After the presort, the number of bivalves per each size category should be determined. The test should be started using the minimum size range with a target range from 5 to 10 mm (that is, from 20 to 25 mm, 33 to 38 mm, 36 to 46 mm) that contains the maximum number of individuals.

11.1.1 Unless a non-recirculating flow-through system is used, test organisms should be kept out of water to eliminate exposure to oxygen deficient conditions. If held out of water, they should be kept cool and moist by providing shade (that is, prevent exposure to direct sunlight), ice packs, or moist toweling. If air temperatures are excessively warm, it may be necessary to hold specimens in an ice chest with wet ice. Once the specimens are sorted into size groups, clean water can be added to the containers. Since the density of animals has been reduced by this time and the shells have been cleaned of epiphytic growth, it is easier to maintain them in clean water without adverse effects. Nevertheless, water temperature should be checked regularly and can be maintained within a desired range by placing plastic bags filled with ice in each container. This is particularly critical when working in a laboratory or other facility where room temperatures can exceed temperatures at the deployment site. A rapid rise in temperature of adult organisms which have ripe gametes could induce spawning, which would add another unwanted variable to the test. If in-situ temperature monitors are used during the deployment period, they can also be used during the sorting and holding period to document temperatures.


11.2 *Final Measurements and Distribution*—Once the final size range has been identified by during the coarse length measurements (during the presort), sorting process, shell lengths should be remeasured and weights measured for the first time. After making these measurements, the bivalves are distributed among the cages. An even distribution of test animal size can be achieved by filling the cages in order of increasing or decreasing animal size (5). This applies to both compartmentalized mesh bags which are attached to PVC frames to form a cage, and to rigid compartmentalized trays.

11.2.1 *Setup for Distribution*—If using mesh bags as described in 9.3.2 and 9.3.3, identification tags, made of durable plastic or other inert material, should be attached to the mesh bags, which have been knotted approximately 30 cm from the end. The identification tag should be secured with a plastic cable tie to the bag near the knot. A water-indelible, permanent marker should be used to label tags with both the cage number and the bag number: for example, a label of 2–3, indicates Cage 2, Bag #3. Color-coded beads strung through a plastic cable tie and fastened to the mesh bag can also be used for purposes of identification. This is recommended as a backup in case the identification tag is lost during the study. The colored bead can be used to identify bag number on the PVC frame, which is also labeled with an indelible marker. If using another type of compartmentalized cage, ensure that the cage is clearly identified, and each compartment within the cage is numbered.


11.2.2 Once all mesh bags have been labeled, the bags can be separated into groups according to bag number, with all the bags with a –1 in one group, –2 in another group, and so forth. The distribution process is based on bag number; all bags of a common number are filled at a given time. To initiate the distribution process, gather all bags that have a “–1” on the label; there should be one for each cage number. Attach these bags to the PVC distribution frame in cage number sequence (Fig. 6). If rigid, compartmentalized cages are used, line up the cages so all cages can be filled simultaneously. Place the cage so the Number 1 compartment is in the upper left position.

11.2.3 *Prepare Bivalves for Distribution*— Starting with either the smallest or largest size group, place all bivalves within the size group into a tray or tub containing water. The bivalves need to be maintained in water during the measurement and distribution process, and the water temperature should be maintained as close as possible (approximately  $\pm 5^{\circ}\text{C}$ ) to temperatures at the deployment site. In most temperate latitudes this can be accomplished by placing plastic bags containing wet ice in the tub with the bivalves; the temperature can be monitored with a thermometer that remains in the tray or with in-situ temperature monitors. It is essential that the bivalves be completely submerged and flat on the bottom prior to measurement. Bivalves with air between their valves either float on the surface or sit upright on the bottom. These bivalves should not be used because the air will bias the whole-animal wet-weights (water weighs more than air). Do not use individuals that float, are buoyant at one end, or do not close upon light physical stimulation (that is, agitation of the water around the bivalves or lightly tapping the shell). Bivalves that float contain air which can be released prior to use. Floating individuals can be transferred to a separate container, where, if left undisturbed, they will likely purge the trapped air.

11.2.4 *Initiate Measuring and Distribution*—Under normal conditions (that is, submerged and respiring), the bivalve shells will be slightly agape (approximately 1 mm). Most species will respond to light physical stimulation by tightly closing their shells.



A. Distribution process for mesh bags.



B. Distribution process for rigid compartmentalized cages.

FIG. 6 Distribution Process for Caging Bivalves

Bivalves that do not completely close their shells upon movement or light physical stimulation should be considered unhealthy and should not be used. In addition, bivalves that have broken shells or holes in their shells should not be used.

11.2.4.1 Initiate the distribution process by randomly selecting one specimen from the holding tray, making sure it is alive and shells are tightly closed. Using a paper towel, blot excess water from exterior of the individual ~~and m.~~ Measure its shell length with a caliper and whole-animal wet-weight with an analytical balance. Record these data to a spreadsheet created for summarizing the data. The data can be recorded either electronically or manually. If the data are entered into an electronic spreadsheet, it is recommended that a manual record also be made as a backup in a case of computer failure. Once the specimen is measured and weighed, place it into the first mesh bag (Fig. 6A) on the distribution rack or into the first cell of the first compartmentalized tray (Fig. 6B). For the mesh bags, affix a 10-cm cable tie around the mesh material above this individual. The cable tie should be adjusted so that it is tight enough to prevent the animal from passing through but the cable tie can be moved if necessary. The cable ties or other restricting devices should not be so tight that the mesh is constricted to the point that it does not allow the shell to open. There should be enough slack to allow movement and growth of the individual during the test. Randomly take another specimen from the tray and measure its shell length and weight, recording the data in the spreadsheet. Place this individual into the second mesh bag on the distribution rack or into the first cell in the second compartmentalized tray. For the mesh bag, affix a cable tie. Repeat this process until one individual has been placed into each mesh bag or each compartmentalized tray. Continue adding bivalves, one at a time to either the mesh bags or the compartmentalized trays (Fig. 6A, B). For the mesh bags, complete one “row” before another row is started, until each mesh bag contains the desired number of individuals. When all of the bags on the distribution rack have been filled, remove the bags, knot or cable tie the open end, leaving a tail length of approximately 0.3 m. For the compartmentalized trays, securely affix a mesh cover such that the bivalves can not migrate from one compartment to another. Place the completed bags or compartmentalized trays into a cooler lined with ice and moist paper towels. Repeat the above process until all the mesh bags or compartmentalized trays are filled. This process ensures that each station will have approximately the same number of individuals from the each size group.

#### ~~11.2.5 If using an electronic~~

11.2.5 Electronic spreadsheet, it can be customized to report minimum, maximum and average shell lengths and weights as they are entered. These values can be compared across cages to identify any individuals that are outside the pre- determined size range and to check for a close, even distribution. To ensure statistical similarity among stations, an Analysis of Variance (ANOVA) can be run on both shell length and whole-animal wet-weight data. The data can be analyzed by cage and by station (that is, pooled cages) if more than one cage is to be deployed at each of the test stations. If statistically significant differences are found, the test animals ~~should~~ can be redistributed to eliminate this difference. For mesh bags, redistribution may require cutting a small opening in the mesh to replace the outlier. The opening can be secured with a plastic cable tie after inserting a replacement individual. For rigid compartmentalized trays, redistribution will require removing the mesh top.

11.2.6 *Mesh Bags Post-distribution Activities*—Once the bivalves have been distributed to all mesh bags, sort the bags by cage number. If the bags are not to be attached to the PVC cages until ~~a later time,~~ later, secure these bags together with a large plastic cable tie. Then, transport bivalves to a pre-identified holding area in the laboratory or field with uncontaminated ~~flowing~~ water. Hold in the water overnight, or until ready to attach bags to the PVC cages.

11.2.7 *Rigid Compartmentalized Trays Post-distribution Activities*—Once the bivalves have been distributed to all the compartmentalized trays, transport bivalves to a pre-identified holding area in the laboratory or field with uncontaminated ~~flowing~~ water. Hold in the water overnight, or until ready to deploy.

11.2.8 *Measuring and Distributing the Baseline (Beginning-of-Test) Tissue Chemistry Individuals*—Bivalves to be used for baseline tissue chemistry should be identified and separated during the measurement and distribution of test specimens to ensure similar sizes. This can be accomplished by assigning a cage for each baseline tissue chemistry sample. However, instead of distributing the bivalves for baseline tissue chemistry to mesh bags, they should be distributed to rigid, compartmentalized cages. The use of compartmentalized trays for the tissue chemistry specimens eliminates the need to remove individuals from the mesh bags once the distribution process is completed. The individuals should be placed into the compartmentalized trays in order, that is, with the first individual measured placed into compartment #1, the second into compartment #2, etc. At the end of the distribution process, the tissues should be removed for chemical analysis according to the procedures in 11.11.

11.3 *Attachment of Mesh Bags to PVC Frames*—A set of mesh bags attached to any more rigid frame, such as PVC-f or heavy protective mesh, constitutes a cage. Attach the mesh bags to the PVC frame by knotting the tail ends of the mesh directly to the PVC. If there is insufficient material to make a secure knot, use 6 in. (approximately 15 cm) cable ties to firmly attach mesh to the PVC frame. Allow a little slack in the mesh bag during attachment; the mesh should not be stretched so tightly that it restricts bivalve movement. If a temperature recording device is used, it should be attached to the frame at this time. If predators are of concern, wrap the PVC frame with a heavy duty plastic protective mesh, with a mesh size appropriate to deter predators of concern (for example, approximately 1 to 2.5-cm mesh size).

11.3.1 It may be necessary to adjust the space between bivalves in the mesh bags so that the mesh bags can be attached to the PVC frame without being too taut or too loose. During the attachment process, slide the cable ties as necessary to increase or decrease the space between individuals without compromising the space available for each individual (that is, do not decrease the space between animals so that there is insufficient space for them to open their valves during respiration).

11.4 *Deployment*—Deploy caged bivalves at stations in accordance with the procedures in 9.4.

~~11.5 Retrieval and End-of-Test Measurements—At each station, after retrieving the caged bivalves remove any foreign—~~  
Foreign material, if present can be removed, by dipping into the water. The exterior of the shells and the mesh bags should can  
be wiped with paper towels if a sheen to remove fouling or other coating is p not resmoved by dipping. If the bags of bivalves  
 are removed from the PVC frames, a separate ice chest lined with wet ice and moist paper towels should be used to transport the  
 bivalves to the processing site. ~~Otherwise, cover the bivalve cages with~~ place a tarp or other protective covering under and over  
the bivalve cages during transportation to prevent contact with contaminated surfaces ans desiccation.

11.5.1 Guide E 1688 recommends a 24-h gut purging period for deposit-feeding bivalves tested under laboratory conditions. Guide E 1688 also gives suggestions when not to purge the gut. It is up to the investigator to determine if the experimental design requires gut purging. However, in the field a 24-h gut purging may not be convenient or necessary due to time and cost constraints. A number of studies have been conducted in which gut purging ranged from 0 to 24 h, such as site-specific bioassays (3, 15) and national, state, and regional mussel watch monitoring programs (45, 46, 205). Gut purging can be accomplished by suspending the caged bivalves at the reference station(s), or at another location known to be relatively free of contamination when compared to the test stations, for appropriate time periods.

~~11.5.2 At the processing site, bivalves~~

11.5.2 Bivalves from all bags constituting a cage or all bivalves in one compartmentalized tray should be processed together. It is critical to retain the order of bivalves during the end-of-test measurements. For mesh bags, it is recommended to remove the bivalves from the mesh bags and place them in a rigid compartmentalized tray to maintain order and facilitate end-of-test measurements. Start with Bag-1 from a given cage. Starting at the end of the bag with the plastic label, remove the bivalves and place them, in sequential order, into a compartmentalized plastic tray. Trays with holes to allow water circulation are recommended. After all individuals from Bag-1 have been transferred to the compartmentalized tray, repeat the process with the remaining bags, maintaining bag sequence (that is, process Bag-2, then Bag-3, and so forth). If a dead or missing individual is encountered, leave its corresponding compartment in the tray empty, or place a marker (that is, a bead or other device) in the compartment. This will ensure the order of individuals is maintained. Depending on the number of bivalves used, it may be necessary to use more than one compartmentalized tray to hold all the bivalves from a given cage. For bivalves deployed in rigid compartmentalized trays, remove the mesh cover. Approximately 5 to 10 min before initiating length and weight measurements, set the tray(s) into a tub or larger tray containing clean water. The water can be collected from the reference station, the holding facility, or another source of clean water. Upon placement of the compartmentalized tray containing bivalves into the tub of water, some individuals may “float,” indicating air trapped between their valves. It is essential that the bivalves do not float prior to making the weight measurements since the presence of air will compromise the whole-animal weight measurements. It may be necessary to leave the bivalves undisturbed for approximately 5 to 10 min in the water in order for them to open their valves and release the trapped air. Once the air has been released, the bivalves can be taken from the tray and measured.

11.5.3 Starting with the bivalve in the Number 1 compartment, begin the shell length and whole-animal wet-weight measurements. Make sure the individual is alive and the shells are tightly closed before removing it from the compartmentalized tray. The end-of-test measurement procedures are similar to those in 11.2.4: blot the excess water from the exterior of the individual’s shells, measure shell length along the longest axis, and obtain a whole-animal wet-weight measurement. Record the data, either electronically or manually onto spreadsheets. After the individual is measured, place it into a separate compartmentalized tray in the Number 1 compartment. Do not put these compartmentalized trays in larger tubs containing water as it is not necessary to keep bivalves in water once the growth measurements are made. Continue the end-of-test measurements, measuring one individual at a time and retain the order of individuals. For dead or missing individuals, transfer the marker from one compartmentalized tray to the other and enter a “M” or “D” into the spreadsheet to indicate “missing” or “dead.” After all bivalves in a cage are measured and weighed, begin the tissue removal process as described in accordance with 11.7.

11.6 *Analysis of Tissues for Background Contamination* —For in-situ field studies with an exposure component, the initial or background concentration of chemicals in tissues of the test organisms should be analyzed for the chemicals of concern. It may also be necessary to characterize tissue chemistry of the source population well in advance of initiating a test to confirm their appropriateness as an uncontaminated source. However, exposure studies may be conducted without prior chemical analysis of tissues if the bivalves are collected from an area that is monitored for chemical contamination and known to be free of toxicants, or if the tissues of those bivalves have been monitored regularly as in culture facilities. Bivalves collected from unmonitored or potentially contaminated areas should not be used in field bioassays unless caution. It is recommended that their tissues are be  
analyzed for baseline, beginning-of-test chemical concentrations. Bivalves collected from potentially contaminated areas should  
concentrations to confirm they are not be used in field bioassays unless the experimental design specifically requires use of that  
population. contaminated. For in-situ studies that only assess effects, it is not necessary to analyze tissues for baseline, beginning-of-test chemical concentrations. However, as beginning- and end-of-test tissue chemistry data can aid in the interpretation of effects data, these analyses are recommended.

11.7 *Collection and Preparation of Bivalve Tissues for Chemical Analysis*—All equipment used for tissue extraction/collection should be of corrosion resistant stainless steel, anodized aluminum, or borosilicate glass. If corrosion resistant stainless steel is unavailable, use regular stainless steel products, carefully checking before each use for signs of rust, pitting, or corrosion. Do not use if rust, pitting, or corrosion is evident. Before each use, all instruments (that is, cutting board, shucking knife, and weigh pans) should be cleaned in accordance with the minimum following process: wash with a soap-free cleaning solution, hot tap water rinse,

or deionized water rinse. If deemed necessary by the investigator, an acetone, hexane, or 95 % ethanol rinse can follow the last water rinse. Allow the instruments to air dry to remove the potential for adding water to the tissues being collected.

11.7.1 During tissue collection, the order of bivalves should be maintained; tissue weights are recorded by the individual and will be paired with whole-animal wet-weights and other size metrics. Use the compartmentalized trays for holding bivalves prior to shucking, and maintain order after tissues are removed.

11.7.2 If using cutting boards made of a material other than corrosion resistant stainless steel, anodized aluminum, or borosilicate glass, cutting boards should be covered with aluminum foil and cleaned as indicated in 11.7. If gloves are worn during the shucking process, they should be powder free. Wash hands thoroughly with a soap-free cleaning solution, or replace gloves between processing a cage of bivalves.

11.7.3 Tissues are removed in accordance with the following process. Start with the first individual in the compartmentalized tray, work with one individual at a time, and retain the order of individuals. Place bivalve on the cutting board. Slide the knife blade between bivalve shells, severing posterior and anterior adductor muscles. Spread the shells apart to reveal soft tissues. If preparing tissues from clams or oysters, it may be necessary to notch the shell prior to inserting the knife blade between their shells. Use a separate knife designated only for the purpose of shell notching. Be sure that none of this shell material is combined with the soft tissue material. Using the tip of the knife blade, separate tissue from the shell, scraping as much of the adductor muscle from points of attachment as possible. Holding tissues to the shell with the blade of the shucking knife, tip the shell to drain excess liquid (that is, the water that was trapped between the shells during the measurement process).

11.7.4 After complete separation, keep the tissues in the shell and use the shell as a “holding dish” until tissue weights are measured. If the two shell halves should become separated, place one half under the other. Place the shell(s) containing the separated tissue on a tray lined with aluminum foil, keeping bivalves in order and sufficient space between the individuals to prevent the shell of one individual from touching the soft tissue of another. Minimize exposure of bivalve tissue to hands, aluminum foil, and any other surface other than the interior of the specimen’s original shell. Repeat this process until all bivalves constituting a “chemical replicate” are shucked.

11.7.5 Place the weigh pan on the analytical balance; tare balance. Pick up the first specimen, and using the shucking knife blade tip, slide the tissue onto the weigh pan. Allow the balance to stabilize. Record the weight, either electronically or manually. Tare material on balance. Continue adding tissues, one at a time, recording the weights of each individual. Tare after each addition. When all tissues of a “chemical replicate” have been weighed, transfer tissues from the weigh pan to the prepared sample jar by gently sliding them off the foil. Tightly cap the sample jar, affix prepared label, and place the tissue samples in a cool location (that is, ice chest containing gel packs, wet or dry ice, or a freezer) depending on the specifications of the analytical laboratory performing the tissue chemistry analyses. If using aluminum foil to line surfaces, discard foil and clean all sampling equipment in accordance with 11.11 before proceeding to the next sample.

11.8 *Quality Assurance/Quality Control Procedures*—Quality assurance is a program designed to provide accurate and precise results. Included are the selection of proper technical methods, sample collection, selection of limits, and qualifications and training of personnel. Quality control are specific actions required to provide information for the quality assurance program. Included are standardization, calibration, replicates, and control and check samples suitable for statistical estimates of confidence of the data (206).

11.8.1 To ensure that the measuring instruments (that is, calipers and balance) are providing accurate readings, the instrument can be tested by measuring a standard weight or shell length. For the balance, one or more from a series of standard weights (that is, 10, 50, 100 and 200 g) can be applied to the balance at intervals throughout the measurement process. For example, after every 100 measurements made on the balance, a standard 100-g weight can be applied to the balance. If the balance is off by more than 1 % (1 g), the balance should be recalibrated; it may be necessary to reweigh some of the previous individuals depending on the degree of off-calibration. The measurement accuracy of the calipers can be checked by completely closing the device and recording the displayed measurement, which should be 0.000 mm. If the caliper displays a value greater than 0.5 mm, the unit should be re-zeroed. If available, a standard length can be measured to check the accuracy of the calipers. Depending on the degree of off-calibration, it may be necessary to remeasure some of the previous individuals.

11.8.2 Quality assurance/quality control (QA/QC) procedures of bivalve measurements should be used primarily in the development process as practitioners refine their methodology. Once the methods have become routine, it may only be necessary to use the QA/QC procedures on a yearly basis to confirm that no artifacts have inadvertently entered the methodology.

11.8.2.1 One suggested approach for QA/QC procedures for the bivalve measurements is to remeasure and reweigh 5 % of the animals. These QA/QC measurements can be performed during the initial and end-of-test measurement processes. The QA/QC shell length measurements and whole-animal wet-weight measurements outside  $\pm 5$  % of the original measurements may be considered unacceptable error measurements. The remeasuring of animal shell length and weight occurs throughout the measurement process as each series of bags is processed to ensure that all measurements are within the limits defined as acceptable. As an example, suggested limits for *Mytilus* sp. approximately 30 to 40 mm in length are  $1.0 \pm 0.5$ -mm variance in shell length and a  $0.5 \pm 0.25$ -g variance in weight. If the results of the remeasurements fall outside of these limits, it is recommended that the previous batch of 100 individuals be remeasured. The hard copy data sheets can contain a separate row for the QA/QC measurements. To facilitate the process, it is suggested that these QA/QC measurements be made on the last “row” of bivalves to be entered into a series of bags.

11.9 *Sample Containers, Handling, and Preservation*—Pre-cleaned sample containers should be purchased from a supplier or provided by the analytical laboratories. Each jar should be sealed, affixed with a completed label, assigned a unique tag number, and stored under appropriate conditions. Sample labels should be made of self-adhering, waterproof material; an indelible pen should be used to fill out each label. Each sample label should contain the project number, sample identification, preservation technique, analyses, date and time of collection, and initials of the person(s) preparing the sample. A completed sample label should be affixed to each sample container. In addition, a unique numbered tag can be affixed to each sample container. Chain-of-custody forms and tamper-proof tape can be used for projects that are litigation sensitive. The preservation of tissue samples is a function of chemical analytes and methods used by the analytical laboratory. The analytical laboratory should provide guidance on proper handling and preservation of tissue samples. For most analyses, samples should be protected from light and refrigerated at  $4^{\circ} \pm 2^{\circ}\text{C}$  from the time of receipt until they are extracted and analyzed. For some chemical analyses or longer holding times, it may be necessary to freeze the samples at  $-20^{\circ}\text{C}$  (Guide E 1688).

## 12. Ancillary Measurements

12.1 It is recommended that supplementary measurements be made on those factors most likely to affect bioaccumulation and growth. These include factors such as chemicals in water and sediment, temperature, food, dissolved oxygen, pH, salinity, suspended particulate matter, turbidity, and water current velocity. These measurements provide site-specific information that can be used to help explain the measured exposure and associated effects. These measurements are more easily made at shallower depths but may become problematic at the greater depths where caged bivalves have been deployed (for example, 650 m). Nevertheless, it is important to measure the chemicals in water or sediment, and this has been accomplished even at these great depths with discrete samples (for example, (13, 14)). In-situ temperature monitoring devices are available to measure temperatures for periods up to 1 year at depths up to 330 m. Standardized procedures should be used to measure chemicals of concern in water and sediment.

12.2 *Chemical Exposure*—It is highly recommended that the concentration of chemicals in water sediment be determined, in addition to measuring the concentration of chemicals in tissues, to characterize total potential exposure; that is, exposure from both water and sediment. These data can be used to establish relationships between chemicals in various environmental compartments as a form of field validation or to establish first-order approximations of these relationships that are validated in the laboratory. Results of the bivalve field bioassay can be used to predict bioaccumulation and biological effects likely to occur in other aquatic organisms under comparable field conditions. While this may only represent first order approximations in some cases, equilibrium partitioning theory, quantitative structure activity relationships, and critical body residue theory suggest that tissue burdens of chemicals associated with adverse effects may be similar across species (6, 7).

12.3 *Temperature*—Marine and freshwater species should be selected to match the site-specific temperatures in the area of concern. Ideally, if species are naturally found in the area or have been found in the area in the past, it is a good indication that temperature tolerances are appropriate. Since temperature could influence bioaccumulation and growth (2, ~~55~~ 57), it is important to monitor temperature during the course of the test using in-situ temperature monitors.

12.4 *Food*—As with temperature, if indigenous populations of the bivalves of choice are found in the area of concern, it is a good indication that there is adequate food to support caged bivalves in the area. Since food could also influence bioaccumulation and growth (2, ~~55~~ 57), it may be helpful to measure parameters such as chlorophyll-a, particulate or total organic carbon, and suspended solids during the course of the test.

## 13. Acceptability of Test

13.1 An acceptable test should meet both survival and growth criteria because the data are intended for different purposes. Survival is a generic indication of overall health and test acceptability. Growth may be a more sensitive indicator of health than survival, and tissue weights should also be monitored. There should be no significant loss in tissue weight during the exposure period. This is necessary for meaningful bioaccumulation potential even if significant growth does not occur. It may be necessary to use different criteria for exposure and effects end points. If survivors have not lost significant tissue mass, a survival criterion of  $>45\%$  may be acceptable to interpret the bioaccumulation data. Conversely, since more individuals may be necessary to interpret effects end points like growth, a 45 % survival criterion may not be acceptable.

13.2 There should not be significant loss in either end-of-test tissue weights or whole-animal wet-weights when compared to measurements made at the beginning of the test, particularly at reference station(s), if they are used. If tissue weights decrease by more than 20 %, this could provide valuable effects information. However, this loss in tissue weight could be accompanied by loss in chemicals and represents a biased estimate of potential chemical bioavailability. Therefore, results may be considered unacceptable for interpreting the environmental significance of bioaccumulation if (1) the end-of-test tissue weights are more than 20 % lower than the beginning-of-test estimates, and (2) the end-of-test tissue weights are significantly less ( $\alpha = 0.05$ ) than the beginning-of-test estimates.

## 14. Report

14.1 A record of the results of an acceptable caged bivalve exposure and effects test should include the following information:

14.1.1 Names of test and investigator(s), name and location of laboratory, and dates of initiation and termination of the test;

14.1.2 Source of test animals, scientific name and how verified, initial whole-animal wet-weights, shell lengths, and estimates

of tissue weights as well as end-of-test percent survival, whole animal wet-weights, shell lengths, and estimates of tissue weights. Means, ranges, and standard deviations of all measurements.

14.1.3 Description of the experimental design and cages, including any attached instrumentation and predator deterring devices, water depth and depth of cages at time of deployment, the number of animals per station, station coordinates, and any other outstanding features of the area to assist in station identification.

14.1.4 Averages and ranges of the acclimation temperature during the measurement and distribution process as well as the time spent out of water while in transit to the measurement location at the beginning of the test and while in transit to the deployment locations at the beginning and end of the test.

14.1.5 Reproductive state of the test animals including degree of gonad development by observation, mass, or volume measurements. Note whether bivalves spawned during the beginning-of-test or end-of-test measurements.

14.1.6 A table of data on concentrations of chemicals in water, sediment, and tissues (including percent lipids and percent moisture in tissues) should be included with sufficient detail to allow independent statistical analyses. The table should also include analytical methods and laboratory qualifiers.

14.1.7 A table of survival, effects, and tissue chemistry data (including percent lipids and percent moisture of the tissues) in sufficient detail to allow independent statistical analyses.

14.1.8 Anything unusual about the test, any deviation from these procedures, and any other relevant information.

14.1.9 Published reports should contain enough information to clearly identify the methodology used and the quality of the results.

## 15. Keywords

15.1 bioaccumulation; bivalve; exposure effects; field bioassay; growth; in-situ

## REFERENCES

- (1) Bayne, B. L., Brown, D. A., Burns, K., Dixon, D. R., Ivanovici, A., Livingstone, D. R., Lowe, D. M., Moore, N. M., Stebbing, A. R. D., and Widdows, J., *The Effects of Stress and Pollution on Marine Animals*, Praeger Special Studies, Praeger Scientific, New York, 1985.
- (2) Widdows, J., and Donkin, P., "Mussels and Environmental Contaminants: Bioaccumulation and Physiological Aspects," *The Mussel Mytilus: Ecology, Physiology, Genetics and Culture*, E. Gosling, ed., Elsevier Science Publishers, Amsterdam, 1992, pp. 383-424.
- (3) Salazar, M. H., and Salazar, S. M., "Mussels As Bioindicators: Effects of TBT on Survival, Bioaccumulation and Growth Under Natural Conditions," *Tributyltin: Environmental Fate and Effects*, M. A. Champ, and P. F. Seligman, eds., Chapman and Hall, London, 1996, pp. 305-330.
- (4) Salazar, M. H., and Salazar, S. M., "Assessing Site-Specific Effects of TBT Contamination With Mussel Growth Rates," *Mar. Env. Res.*, Vol 32, No. 1-4, 1991, pp. 131-150.
- (5) Salazar, M. H., and Salazar, S. M., "In Situ Bioassays Using Transplanted Mussels: I. Estimating Chemical Exposure and Bioeffects With Bioaccumulation and Growth," *Environmental Toxicology and Risk Assessment - Third Volume*, J. S. Hughes, G. R. Biddinger, and E. Mones, eds., American Society for Testing and Materials, Philadelphia, 1995, pp. 216-241.
- (6) McCarty, L. S., "Toxicant Body Residues: Implications for Aquatic Bioassays With Some Organic Chemicals," *Aquatic Toxicology and Risk Assessment*, M. A. Mayes, and M. G. Barron eds., American Society for Testing and Materials, Philadelphia, 1991, pp. 183-192.
- (7) McCarty, L. S., and Mackay, D., "Enhancing Ecotoxicological Modeling and Assessment," *Environ. Sci. Technol.*, Vol 27, No. 9, 1993, pp. 1719-1728.
- (8) US ACOE, 1999, Environmental Residue Effects Database Home Page. (Web Page), Available at <http://www.wes.army.mil/el/ered/index.html#misc>.
- (9) Jarvinen, A. W., and Ankley, G. T., *Linkage of Effects to Tissue Residues: Development of a Comprehensive Database for Aquatic Organisms Exposed to Inorganic and Organic Chemicals*, SETAC, Pensacola, FL, 1999.
- (10) US EPA, 1998a, *Guidelines for Ecological Risk Assessment - Final Risk Assessment Forum*, US Environmental Protection Agency, Washington, DC EPA/630/R-95/002F.
- (11) Luoma, S. N., "Prediction of Metal Toxicity in Nature From Bioassays: Limitations and Research Needs," *Metal Speciation and Bioavailability*, A. Tessier, and D. Turner, eds., John Wiley & Sons, Chichester, 1995, pp. 609-659.
- (12) Luoma, S. N., and Fisher, N., "Uncertainties in Assessing Contaminant Exposure From Sediments: Bioavailability," *Ecological Risk Assessment of Contaminated Sediments*, C. G. Ingersoll, T. Dillon, and G. Biddinger, eds., SETAC Press, Pensacola, FL, 1997.
- (13) Forlin, L., Baden, S. P., Eriksson, S., Granmo, A., Lindesjoo, E., Magnusson, K., Ekelund, R., Esselin, A., and Sturve, J., "Effects of Contaminants in Roundnose Grenadier (*Corphaenoides rupestris*) and Norway Lobster (*Nephrops norvegicus*) and Contaminant Levels in Mussels (*Mytilus edulis*) in the Skagerrak and Kattegat Compared to the Faroe Islands," *J. Sea Res.*, Vol 35, No. 1-3, 1996, pp. 209-222.
- (14) Forlin, L., Livingstone, D. R., Magnusson, K., Peters, L. D., Sole, M., Sjolone, A., and Granmo, A., "Molecular Investigations into Pollutant Impact on Roundnose Grenadier (*C. rupestris*) and Transplanted Common Mussel (*Mytilus edulis*) in Skagerrak, the North Sea," *Mar. Environ. Res.*, Vol 42, No. 1-4, 1996, pp. 209-212.
- (15) Salazar, M. H., and Salazar, S. M., "Using Bioaccumulation and Growth in Caged Intertidal Oysters to Assess Oil Exposure and Effects in Delaware Bay", *Proceedings of the Twentieth Arctic and Marine Oilspill Program (AMOP) Technical Seminar*, Vancouver, British Columbia, Environment Canada, 1997, pp. 661-675.
- (16) Di Toro, D. M., McGrath, J. A., and Hansen, D. J., In Review, "Technical Basis for Narcosis Chemicals and PAH Criteria. I. Water and Tissue," *Submitted to Environ. Toxic. Chem.*
- (17) Bridges, T. S., Moore, D. W., Landrum, P., Neff, J., and Cura, J., "Summary of a Workshop on Interpreting Bioaccumulation Data Collected During Regulatory Evaluations of Dredged Material. Miscellaneous Paper D-96-July 1996," Denver, CO, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS, 1996.
- (18) Salazar, M. H., and Salazar, S. M., "Using Caged Bivalves As Part of an Exposure-Dose-Response Triad to Support an Integrated Risk Assessment

- Strategy,” *Proceedings, Ecological Risk Assessment: A Meeting of Policy and Science, SETAC Special Publication*, A. de Peyster, and K. Day, eds., SETAC Press, 1998, pp. 167-192.
- (19) Bright, D. A., “Tissue Variability in the Infaunal Bivalve *Axinopsida serricata* (Lucinacea: Thyasiridae) Exposed to a Marine Mine Tailings Discharge; and Associated Population Effects,” Victoria, B.C., University of Victoria, 1991.
  - (20) Blaise, C., Gagné, F., Pellerin, J., and Hansen, P. D., “Determination of Vitellogenin-Like Properties in *Mya arenaria* Hemolymph (Saguenay Fjord, Canada): a Potential Biomarker for Endocrine Disruption,” *Environ. Toxicol.*, Vol 14, No. 5, 1999, pp. 455-465.
  - (21) Widdows, J., Phelps, D. K., and Galloway, W., “Measurement of Physiological Condition of Mussels Transplanted Along a Pollution Gradient in Narragansett Bay,” *Mar. Environ. Res.*, Vol 4, 1981, pp. 181-194.
  - (22) Bruner, K. A., Fisher, S. W., and Landrum, P. F., “The Role of the Zebra Mussel, *Dreissena polymorpha*, in Contaminant Cycling: I. The Effect of Body Size and Lipid Content on the Bioconcentration of PCBs and PAHs,” *J. Great Lakes Res.*, Vol 20, No. 4, 1994, pp. 725-734.
  - (23) Bruner, K. A., Fisher, S. W., and Landrum, P. F., “The Role of the Zebra Mussel, *Dreissena polymorpha*, in Contaminant Cycling: II. Zebra Mussel Contaminant Accumulation from Algae and Suspended Particles, and Transfer to the Benthic Invertebrate, *Gammarus fasciatus*,” *J. Great Lakes Res.* Vol 20, No. 4, 1994, pp. 735-750.
  - (24) Harvey, R. W., and Luoma, S. N., “Separation of Solute and Particulate Vectors of Heavy Metal Uptake in Controlled Suspension-Feeding Experiments with *Macoma balthica*,” *Hydrobiologia*, Vol 121, 1985, pp. 97-102.
  - (25) Lee, B. G. and Luoma, S. N., “Influence of Microalgal Biomass on Absorption Efficiency of Cd, Cr, and Zn by Two Bivalves from San Francisco Bay,” *Limnol. Oceanogr.*, Vol 43, No. 7, 1998, pp. 1455-1466.
  - (26) Luoma, S. N., “The Developing Framework of Marine Ecotoxicology: Pollutants as a Variable in Marine Ecosystems,” *J. Exp. Mar. Biol. Ecol.* Vol 200, 1996, pp. 29-55.
  - (27) Luoma, S. N., Johns, C., Fisher, N. S., Steinberg, N. A., Oremland, R. S., and Reinfelder, J. R., “Determination of Selenium Bioavailability to a Benthic Bivalve from Particulate and Solute Pathways,” *Environ. Sci. Technol.* Vol 26, 1992, pp. 485-491.
  - (28) Hyland, J.L., Van Dolah, R.F., and Snoots, T. R., “Predicting Stress in Benthic Communities of Southeastern U.S. Estuaries in Relation to Chemical Contamination in Sediments,” *Environ. Toxicol. Chem.* 18(11), 1998, pp. 2557-2564.
  - (29) Hyland, J.L., Van Dolah, R.F., and Snoots, T. R., “Predicting Stress in Benthic Communities of Southeastern U.S. Estuaries in Relation to Chemical Contamination in Sediments,” *Environ. Toxicol. Chem.* 18(11), 1999, pp. 2557-2564.
  - (30) Doherty, F. G., Cherry, D. S., and Cairns, J. Jr., “Valve Closure Responses of the Asiatic Clam *Corbicula fluminea* Exposed to Cadmium and Zinc,” *Hydrobiologia*, Vol 153, No. 2, 1987, pp. 159-168.
  - (31) Redpath, K.J. and J. Davenport 1988. The Effect of Copper, Zinc and Cadmium on the Pumping Rate of *Mytilus edulis* L. *Aquat. Toxicol.* 13(3):217-226.
  - (32) Trueman, E. R., and Akberali, H. B., “Responses of an Estuarine Bivalve, *Scrobicularia plana* (Tellinacea) to Stress,” *Malacologia*, Vol 21, No. 1-2, 1981, pp. 15-21.
  - (33) Waller, W. T., Acevedo, M. F., Morgan, E. L., Dickson, K. L., Kennedy, J. H., Ammann, L. P., Allen, J. H., and Keating, P. R., “Biological and Chemical Testing in Storm Water”, *Stormwater NPDES Related Monitoring Needs, Proceedings of an Engineering Foundation Conference*, Mount Crested Butte, CO, 1994, pp. 177-193.
  - (34) Keller, A. E., and Zam, S. G., “The Acute Toxicity of Selected Metals to the Freshwater Mussel, *Anodonta imbecilis*,” *Environ. Toxicol. Chem.* Vol 10, 1991, pp. 539-546.
  - (35) Moore, J. W., Somers, J. D., Fritz, D. L., Smiley, K. L., Goski, B., and Blumhagen, K., “Toxicity of Municipal Wastewater to Two Species of Fish, the Cladoceran *Daphnia magna*, and the Mollusc, *Anodonta grandis*,” *Proceedings of the Nineteenth Annual Aquatic Toxicity Workshop*, E. G. Baddaloo, S. Ramamoorthy, and J. W. Moore, eds., Edmonton, Alberta, Canadian Technical Report of Fisheries and Aquatic Sciences, 1992, pp. 145-155.
  - (36) McKinney, A. D., and Wade, D. C., “Comparative Response of *Ceriodaphnia dubia* and Juvenile *Anodonta imbecillis* to Pulp and Paper Mill Effluents Discharged to the Tennessee River and Its Tributaries,” *Environ. Toxicol. Chem.*, Vol 15, No. 4, 1996, pp. 514-517.
  - (37) US EPA, 1998 *Update for Ambient Water Quality Criteria for Ammonia*, U.S. Environmental Protection Agency, Office of Water, August 1998.
  - (38) Burgess, R. M., and Morrison, G. E., “A Short-Exposure, Sublethal, Sediment Toxicity Test Using the Marine Bivalve *Mulinia lateralis*: Statistical Design and Comparative Sensitivity,” *Environ. Toxicol. and Chem.* Vol 13, No. 4, 1994, pp. 571-580.
  - (39) Van Dolah, R. F., Maier, P. P., Jones, J. D., Ringwood, A. H., Keppler, C. J., Conners, D. E., Fulton, M. H., Scott, G. I., Chung, K., Chandler, G. T., Lee, R. F., Snell, T. W., and Cecchine, G. A., “A Comparison of Sediment Bioassay Protocols Involving Contaminant Mixtures (Abstract Only),” *SETAC 19th Annual Meeting. The Natural Connection: Environmental Integrity and Human Health*, Charlotte, NC, Society of Environmental Toxicology and Chemistry, 1998.
  - (40) Black, M. C., and Belin, J. I., “Evaluating Sublethal Indicators of Stress in Asiatic Clams (*Corbicula fluminea*) Caged in an Urban Stream,” *Environmental Toxicology and Risk Assessment, Seventh Volume*, E. E. Little, B. M. Greenberg, and A. J. DeLonay, eds., ASTM STP 1333, Fredericksburg, VA, 1998, pp. 76-91.
  - (41) Black, M. C., Ferrell, J. R., Horning, R. C., and Martin, L. K. Jr., “DNA Strand Breakage in Freshwater Mussels (*Anodonta grandis*) Exposed to Lead in the Laboratory and the Field,” *Environ. Toxicol. Chem.*, Vol 15, No. 5, 1996, pp. 802-808.
  - (42) Couillard, Y., Campbell, P. G. C., Pellerin-Massicotte, J., and Auclair, J. C., “Field Transplantation of a Freshwater Bivalve, *Pyganodon grandis*, Across a Metal Contamination Gradient. I. Temporal Changes in Metallothionein and Metal (Cd, Cu, and Zn) Concentrations in Soft Tissues,” *Can. J. Fish. Aquat. Sci.* Vol 52: 1995, pp. 690-702.
  - (43) Couillard, Y., Campbell, P. G. C., Pellerin-Massicotte, J., and Auclair, J. C., “Field Transplantation of a Freshwater Bivalve, *Pyganodon grandis*, Across a Metal Contamination Gradient. II. Metallothionein Response to Cd and Zn Exposure, Evidence for Cytotoxicity, and Links to Effects at Higher Levels of Biological Organization,” *Can. J. Fish. Aquat. Sci.* Vol 52, 1995, pp. 703-715.
  - (44) Couillard, Y., Parisot, Y., and Campbell, P. G. C., “Metallothionein Concentrations and Cytosolic Distributions in Indigenous Freshwater Benthic Invertebrates From Metal-Contaminated Environments,” *Proceedings of the 21st Annual Aquatic Toxicity Workshop*, G. F. Westlake, J. L. Parrott, and A. J. Niimi, eds., Sarnia, Ontario, 1995, pp. 126.
  - (45) O’Connor, T. P., Cantillo, A. Y., and Lauenstein, G.G., “Chapter 2 - Monitoring of Temporal Trends in Chemical Contamination by the NOAA National Status and Trends Mussel Watch Project,” *Biomonitoring of Coastal Waters and Estuaries*, K. J. M., Kramer, ed., CRC Press, Inc., Boca Raton, FL, 1994, pp. 29-50.
  - (46) Martin, M., and Severeid, R., “Mussel Watch Monitoring for the Assessment of Trace Toxic Constituents in California Marine Waters,” *Concepts*

- in *Marine Pollution Measurements*, H. H. White, ed., Maryland Sea Grant Publication, University of Maryland, College Park, 1984, pp. 291-323.
- (47) State of California, "Toxic Substances Monitoring Program: Ten Year Summary Report 1978-1987. 90-1WQ," Water Resources Control Board, State of California, 1990.
  - (48) Naimo, T. J., "A Review of the Effects of Heavy Metals on Freshwater Mussels," *Ecotoxicology*, Vol 4, 1995, pp. 341-362.
  - (49) Mersch, J., and Pihan, J. C., "Simultaneous Assessment of Environmental Impact on Condition and Trace Metal Availability in Zebra Mussels *Dreissena polymorpha* Transplanted into the Wiltz River, Luxembourg. Comparison With the Aquatic Moss," *Arch. Env. Contam. Toxicol.*, Vol 25, 1993, pp. 353-364.
  - (50) Mersch, J., Wagner, P., and Pihan, J. C., "Copper in Indigenous and Transplanted Zebra Mussels in Relation to Changing Water Concentrations and Body Weight," *Environ. Toxicol. Chem.*, Vol 15, No. 6, 1996, pp. 886-893.
  - (51) Morrison, H. A., Lazar, R., and Haffner, G. D., "In Situ Bioaccumulation of Polychlorinated Polychlorinated Biphenyls by Zebra Mussels," *Proceedings of the 21st Annual Aquatic Toxicity Workshop*, G. F. Westlake, J. L. Parrott, and A. J. Niimi, eds., Sarnia, Ontario, 1995, pp. 128.
  - (52) Reincke, H., "Biological Effect Monitoring in the River Elbe Using the Zebra Mussel *Dreissena polymorpha*," *Limnologie Aktuell*, Vol 4, 1992, pp. 185-196.
  - (53) Hilbish, T. J., "Growth Trajectories of Shell and Soft Tissue in Bivalves: Seasonal Variation in *Mytilus Edulis* L.," *J. Exp. Mar. Biol. Ecol.* Vol 96, 1986, pp. 103-113.
  - (54) Lewis, D. E., and Cerrato, R. M., "Growth Uncoupling and the Relationship Between Shell Growth and Metabolism in the Soft Shell Clam *Mya arenaria*," *Mar. Ecol. Prog. Ser.*, Vol 158, 1997, pp. 177-189.
  - (55) Bayne, B. L., *Marine Mussels: Their Ecology and Physiology*, Cambridge University Press, Cambridge, 1976.
  - (56) Salazar, M. H., and Kenis, P. R., "Evaluation of Sewage Toxicity - a Comparison of Bioassay Techniques," *Naval Undersea Center Tech. Note 1251*, 1973.
  - (57) Martin, J. M., Piltz, F. M., and Reish, D. J., "Studies on the *Mytilus edulis* Community in Alamitos Bay, California. The Effects of Heavy Metals on Byssal Thread Production," *Veliger*, Vol 18, No. 2, 1975, pp. 183-188.
  - (58) Van Winkle, W. Jr., "Effect of Environmental Factors on Byssal Thread Formation," *Mar. Biol.*, Vol 7, 1970, pp. 143-148.
  - (59) Huggett, R. J., Kimerle, R. A., Mehrle, P. M. Jr., and others, *Biomarkers: Biochemical, Physiological, and Histological Markers of Anthropogenic Stress*, Lewis Publishers, Boca Raton, FL, 1992.
  - (60) McCarthy, J. F., and Shugart, L. R., *Biomarkers of Environmental Contamination*, Lewis Publishers, Boca Raton, FL, 1990.
  - (61) Koepf, S. J., Santoro, E. D., Zimmer, R., and Nadeau, J., "Bioaccumulation of Hg, Cd, and Pb in *Mytilus edulis* Transplanted to a Dredged- Material Dumpsite," *Oceanic Processes in Marine Pollution, Volume 1, Biological Processes and Wastes in the Ocean*, J. M. Capuzzo, and D. R. Kester, eds., Krieger Publishing, Malabar, FL, 1987.
  - (62) Gentile, J. H., Bierman, V. J. Jr., Paul, J. F., Walker, H. A., and Miller, D. C., "A Hazard Assessment Research Strategy for Ocean Disposal," *Oceanic Processes in Marine Pollution, Volume 1, Biological Processes and Wastes in the Ocean*, J. M. Capuzzo, and D. R. Kester, eds., Krieger Publishing, Malabar, FL, 1987, pp. 199-212.
  - (63) Paul, J. F., Bierman, V. J. Jr., Davis, W. R., Hoffman, G. L., Munns, W. R., Pesch, C. E., Rogerson, P. F., and Schimmel, S. C., "The Application of a Hazard Assessment Research Strategy to the Ocean Disposal of a Dredged Material: Exposure Assessment Component," *Oceanic Processes in Marine Pollution. Volume 4, Scientific Monitoring Strategies for Ocean Waste Disposal*, D. W. Hood, A. Schoener, and P. K. Park, eds., Krieger Publishing, Malabar FL, 1989, pp. 123-135.
  - (64) Phelps, D. K., Johns, D. M., Scott, K. J., Galloway, W. B., Reynolds, B. H., Nelson, W. G., Rosen, J. S., Black, D., Lake, J. L., Gutjahr-Gobell, R., Lussier, S., Mueller, C., Redmond, M., Schauer, P. S., Yevich, C. A., Yevich, P. P., Zaroogian, G. E., and Heltshe, J., "The Application of a Hazard Assessment Research Strategy to the Ocean Disposal of a Dredged Material: Effects Assessment and Monitoring Component.," *Oceanic Processes in Marine Pollution; Volume 4: Scientific Monitoring Strategies for Ocean Waste Disposal*, D. W. Hood, A. Schoener, and P. K. Park, eds., Krieger Publishing, Malabar, FL, 1989, pp. 137-145.
  - (65) Nelson, W. G., and Hansen, D. J., "Development and Use of Site-Specific Chemical and Biological Criteria for Assessing New Bedford Harbor Pilot Dredging Project," *Environmental Manag.*, Vol 15, No. 1, 1991, pp. 105-112.
  - (66) Young, D. R., "Chlorinated Hydrocarbon Contaminants in the Southern California and New York Bights," *Ecological Stress and the New York Bight: Science and Management*, G. F. Mayer, ed., Estuarine Research Federation, Columbia, SC, 1982, pp. 263-276.
  - (67) National Oceanic and Atmospheric Administration and EVS Consultants, "Baseline Clam Monitoring Study, Cannelton Industries Site, Sault Ste. Marie, Michigan, Summer 1997. September 1998," NOAA Technical Memorandum NOS ORCA 136.
  - (68) Matta, M. B., Salazar, S., Mill, L., Gray, G., Linse, J., Peronard, P., and Francendese, L., "Final Draft. LCP Chemical Site, Monitoring Study Data Report," National Oceanic and Atmospheric Administration. U.S. Environmental Protection Agency, Region IV, May 27, 1998.
  - (69) Herve, S., Prest, H. F., Heinonen, P., Hyorilainen, T., Koistinen, J., and Paasivirta, J., "Lipid-Filled Semipermeable Membrane Devices and Mussels as Samplers of Organochlorine Compounds in Lake Water," *Environ. Sci. Pollut. Res.*, 2(1), 1995, pp. 24-30.
  - (70) Peven, C. S., Uhler, A. D., and Querzoli, F. J., X "Caged Mussels and Semipermeable Membrane Devices As Indicators of Organic Contaminant Uptake in Dorchester and Duxbury Bays, Massachusetts," *Environ. Toxicol. Chem.*, Vol 15, No. 2, 1996, pp. 144-149.
  - (71) Prest, H. F., Jarman, W. M., Burns, S. A., Weismuller, T., Martin, M., and Huckins, J. N., "Passive Water Sampling Via Semipermeable Membrane Devices (SPMDs) in Concert With Bivalves in the Sacramento/San Joaquin River Delta," *Chemosphere*, Vol 25, No. 12, 1992, pp. 1811-1823.
  - (72) Prest, H. F., Richardson, B. J., Jacobson, L. A., Vedder, J., and Martin, M., "Monitoring Organochlorines With Semi-Permeable Membrane Devices (SPMDs) and Mussels (*Mytilus edulis*) in Corio Bay, Victoria, Australia," *Mar. Pollut. Bull.*, Vol 30, No. 8, 1995, pp. 543-554.
  - (73) Shigenaka, G., and Henry, C. B., "Use of Mussels and Semipermeable Membrane Devices to Assess Bioavailability of Residual Polynuclear Aromatic Hydrocarbons Three Years After the Exxon Valdez Oil Spill," *Third Symposium on Environmental Toxicology and Risk Assessment, ASTM STP 1219, Exxon Valdez Oil Spill: Fate and Effects in Alaskan Waters*, P. G. Wells, J. N. Butler, and J. S. Hughes, eds., American Society for Testing and Materials, Philadelphia, 1995, pp. 239-260.
  - (74) Environment Canada, Interim Sediment Quality Guidelines: Soil and Sediment Quality Section Guidelines Division, Ecosystem Conservation Directorate Evaluation and Interpretation Branch, September 1995.
  - (75) Capuzzo, J. M., Farrington, J. W., Rantamaki, P., Clifford, C. H., Lancaster, B. A., Leavitt, D. F., and Jia, X., "The Relationship Between Lipid Composition and Seasonal Differences in the Distribution of PCBs in *Mytilus edulis* L.," *Mar. Environ. Res.*, Vol 28, 1989, pp. 259-264.
  - (76) McMahon, R. F., Chapter 12 – Ecology of an Invasive Pest Bivalve, *Corbicula*, *The Mollusca*, Volume 6, Ecology, Academic Press, 1983, pp. 505-561.

- (77) Nichols, S. J., Nonindigenous Species Research and Outreach. Life History and Ecological Requirements of the Zebra Mussel - North American Experience Through 1992. <http://www.nsgo.seagrant.org/research/nonindigenous/zmlifehistory.html>, National Sea Grant, 2000.
- (78) Clarke, A. H., "The Freshwater Molluscs of the Canadian Interior Basin," *Malacologia*, 13(1-2), 1973, pp. 1-509.
- (79) McMahon, R. F., Chapter 11 - Mollusca: Bivalvia, *Ecology and Classification of North American Freshwater Invertebrates*, J. H. Thorp and A. P. Covich eds., San Diego, Academic Press, 1991, pp. 315-399.
- (80) Metcalfe-Smith, J. L., Stanton, S. K., Mackie, G. L., and Lane, N. M., "Biodiversity of Freshwater Mussels in the Lower Great Lakes Drainage Basin," *Proceedings, 3rd National Ecological Monitoring and Assessment Network Meeting*, Saskatoon, Saskatchewan, 1997.
- (81) LaSalle, M. W., and de la Cruz, A. A., "Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Gulf of Mexico) – Common Rangia," U.S. Fish Wildl. Serv. Biol. Rep. 82(11.31). U.S. Army Corps of Engineers, TR EL-82-4, 1985, 16 pp.
- (82) Virginia Tech, Fish and Wildlife Information Exchange, Conservation Management Institute, Species Profile. Blacksburg, VA (<http://fwie.fw.vt.edu>), Virginia Tech, 2000.
- (83) Stanley, J. G., and Sellers, M. A., Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Mid-Atlantic) – American Oyster. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.65). U.S. Army Corps of Engineers, TR EL-82-4, 1986, 25 pp.
- (84) Cain, D. J., Studies Using the Clam *Macoma balthica* to Identify Bioavailable Trace Metals in San Francisco Bay. In: The Bioavailability of Toxic Contaminants in the San Francisco Bay-Delta, Proceedings of a Two-Day Seminar Series, October. Report No.: AH1-90-01, 1988.
- (85) Rae, J. G. I., "The Population Dynamics of Two Sympatric Species of *Macoma*," *Veliger* 21(3):1978, pp. 384-399.
- (86) Rae, J. G. I., "Reproduction in Two Sympatric Species of *Macoma* (Bivalvia)," *Biol. Bull.* 155: 1978, pp. 207-219.
- (87) Stanley, J. G., "Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Mid-Atlantic) – Hard Clam," U.S. Fish Wildl. Serv. Biol. Rep. 82(11.41). U.S. Army Corps of Engineers, TR EL-82-4, 1985, 25 pp.
- (88) Newell, C. R. and Hidu, H., "Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (North Atlantic) – Softshell Clam," U.S. Fish Wildl. Serv. Biol. Rep. 82(11.53). U.S. Army Corps of Engineers, TR EL-82-4, 1986, 17 pp.
- (89) Newell, R. I. E., "Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (North and Mid-Atlantic) – Blue Mussel," U.S. Fish Wildl. Serv. Biol. Rep. 82(11.102). U.S. Army Corps of Engineers, TR EL-82-4, 1989, 25 pp.
- (90) Gosling, E., "The Mussel *Mytilus* : Ecology, Physiology, Genetics and Culture," Elsevier, Amsterdam, 1992.
- (91) Skidmore, D., and Chew, K. K., "Mussel Aquaculture in Puget Sound, Seattle, Washington," Washington Sea Grant Publication, 1985.
- (92) Couch, D., and Hassler, T. J., "Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Northwest) – Olympia Oyster," U.S. Fish Wildl. Serv. Biol. Rep. 82(11.124). U.S. Army Corps of Engineers, TR EL-82-4, 1989, 8 pp.
- (93) Shaw, W.N. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Southwest)-Common Littleneck Clam. US Fish Wildl. Serv. Biol. Rep. 82(11.46). US Army Corps of Engineers, TR EL-82-4, 1986, 4pp.
- (94) Short, J. W., and Babcock, M. M., "Prespill and Postspill Concentrations of Hydrocarbons in Mussels and Sediments in Prince William Sound," *Proceedings of the Exxon Valdez Oil Spill Symposium*, S. D. Rice, R. B. Spies, D. A. Wolfe, and others, eds., Anchorage, Alaska, American Fisheries Society, 1996, pp. 149-166.
- (95) Short, J. W., and Heintz, R. A., "Identification of Exxon Valdez Oil in Sediments and Tissues From Prince William Sound and the Northwestern Gulf of Alaska Based on a PAH Weathering Model," *Environ. Sci. Technol.*, Vol 31, No. 8, 1997, pp. 2375-2384.
- (96) Page, D. S., Boehm, P. D., Douglas, G. S., and Bence, A. E., "Identification of Hydrocarbon Sources in the Benthic Sediments of Prince William Sound and the Gulf of Alaska Following the Exxon Valdez Oil Spill," *Exxon Valdez Oil Spill: Fate and Effects in Alaskan Waters*, P. G. Wells, J. N. Butler, and J. S. Hughes, eds., American Society for Testing and Materials, Philadelphia, 1995, pp. 41-83.
- (97) Bence, A. E., and Burns, W. A., "Fingerprinting Hydrocarbons in the Biological Resources of the Exxon Valdez Spill Area," *Exxon Valdez Oil Spill: Fate and Effects in Alaskan Waters*, P. G. Wells, J. N. Butler, and J. S. Hughes, eds., American Society for Testing and Materials, Philadelphia, 1995, pp. 84-140.
- (98) Brumbaugh, W.G., Ingersoll, C.G., Kemble, N.E., May, T.W., Zajicek, J.L., "Chemical Characterization of Sediments and Porewater from the Upper Clark Fork River and Milltown Reservoir, Montana," *Environ. Toxicol. Chem.*, Vol 13, 1994, pp. 1971-1994.
- (99) Anderson, J.W., Jones, J.M., Steinert, S., Sanders, B., Means, J., McMillin, D., Vu, T., and Tukey, R., "Correlation of CYP1A1 Induction, as Measured by the P450 RGS Biomarker Assay, with High Molecular Weight PAHs in Mussels Deployed at Various Sites in San Diego Bay in 1993 and 1995," *Mar. Environ. Res.* Vol 48, 1999, pp. 389-405.
- (100) Zar, J. H., *Biostatistical Analysis*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1974.
- (101) Gilbert, R. O., *Statistical Methods for Environmental Pollution Monitoring*, Van Nostrand Reinhold, New York, 1987.
- (102) Krebs, C. J., *Ecological Methodology*, Harper Collins Publishers, New York, 1989.
- (103) Clark, R. C. Jr., and Finley, J. S., "Uptake and Loss of Petroleum Hydrocarbons by the Mussel *Mytilus edulis*, in Laboratory Experiments," *Fish. Bull.*, Vol 73, No. 3, 1975, pp. 508-515.
- (104) Pittinger, C. A., Buikema, A. L. Jr., Hornor, S. G., and Young, R. W., "Variation in Tissue Burdens of Polycyclic Aromatic Hydrocarbons in Indigenous and Relocated Oysters," *Environ. Toxicol. Chem.* Vol 4, 1985, pp. 379-387.
- (105) Meador, J. P., Stein, J. E., Reichert, W. L., and Varanasi, U., "Bioaccumulation of Polycyclic Aromatic Hydrocarbons by Marine Organisms," *Rev. Environ. Contam. Toxicol.*, Vol 143, 1995, pp. 79-165.
- (106) Naes, K., Bakke, T., and Konieczny, R., "Mobilization of PAH From Polluted Seabed and Uptake in the Blue Mussel (*Mytilus edulis* L.)," *Mar. Freshwater Res.* Vol 46, 1995, pp. 275-285.
- (107) Naes, K., Knutzen, J., and Berglund, L., "Occurrence of PAH in Marine Organisms and Sediments From Smelter Discharge in Norway," *Sci. Tot. Environ.*, Vol 163, 1995, pp. 93-106.
- (108) Calambokidis, J., Mowrer, J., Beug, M. W., and Herman, S. G., "Selective Retention of Polychlorinated Biphenyl Components in the Mussel, *Mytilus edulis*," *Arch. Environ. Contam. Toxicol.* Vol 8, 1979, pp. 299-308.
- (109) Weber, C. I., Report developed jointly by the Office of Research and Development Office of Acid Deposition Environmental Monitoring and Quality Assurance US EPA Cincinnati OH; and the Office of Water US EPA Washington DC, March 1988.
- (110) Dauble, D. D., Daly, D. S., and Abernethy, C. S., "Factors Affecting Growth and Survival of the Asiatic Clam, *Corbicula* sp., Under Controlled Laboratory Conditions," *Aquatic Toxicology and Hazard Assessment: Seventh Symposium, ASTM Special Technical Publication 854*, R. D. Cardwell, R. Purdy, and R. C. Bahner, eds., American Society for Testing and Materials, Philadelphia, 1985, pp. 134-144.
- (111) Negus, C. L., "A Quantitative Study of Growth and Production of Unionid Mussels in the River Thames at Reading," *J. Animal Ecol.*, Vol 35, 1966, pp. 513-532.

- (112) Foe, C., and Knight, A., "Further Studies Evaluating the Freshwater Asiatic Clam, *Corbicula fluminea*, for Monitoring the Sublethal Impact of Point Source Discharges," Prepared for the California Water Quality Control Board, San Francisco Bay Region, (WRCB Standard Agreement No. 2-117-120-0), *Water Science and Engineering Paper No. 4518*, June 1985.
- (113) Foe, C., and Knight, A., "Assessment of the Biological Impact of Point Source Discharges Employing Asiatic Clams," *Arch. Environ. Contam. Toxicol.*, Vol 16, No. 1, 1987, pp. 39-52.
- (114) Muncaster, B. W., Herbert, P. D. N., and Lazar, R., "Biological and Physical Factors Affecting the Body Burden of Organic Contaminants in Freshwater Mussels," *Arch. Environ. Contam. Toxicol.*, Vol 19, 1990, pp. 25-34.
- (115) Salazar, M. H., and Salazar, S. M., In preparation, "Using Mussel Growth Rates to Evaluate Bivalve Biomarkers: Synoptic Testing in San Diego Bay."
- (116) Stephenson, M. D., Coale, S. L., Martin, M., and Martin, J. H., "California Mussel Watch 1979-1980: Trace Metal Concentrations in the California Mussel, *Mytilus californianus*, and the Bay Mussel, *M. edulis*, Along the California Coast and Selected Harbors and Bays. Part 1," Water Quality Monitoring Report 80-8, California State Water Resources Control Board, Sacramento, CA. 1980.
- (117) Stephenson, M. D., Martin, M., and Martin, J. H., "State Mussel Watch - Volume IV: Trace Metal Concentrations in Mussels From Bays of California," California Department of Fish and Game, State Mussel Watch Project, 1980.
- (118) Young, D. R., Heesen, T. C., and McDermott, D. J., "An Offshore Biomonitoring System for Chlorinated Hydrocarbons," *Mar. Pollut. Bull.* Vol 7, No. 8, 1976, pp. 156-160.
- (119) Cain, D. J., and Luoma, S. N., "Copper and Silver Accumulation in Transplanted and Resident Clams (*Macoma balthica*) in South San Francisco Bay," *Mar. Environ. Res.* Vol 15, 1985, pp. 115-135.
- (120) Koch, L., *Surrogates for Threatened and Endangered Species Toxicity Testing*, US Fish and Wildlife Service, Southwest Field Office, Abingdon, VA, 1999.
- (121) Widdows, J., Burns, K. A., Menon, N. R., Page, D. S., and Soria, S., "Measurement of Physiological Energetics (Scope for Growth) and Chemical Contaminants in Mussels (*Arca zebra*) Transplanted Along a Pollution Gradient in Bermuda," *J. Exp. Mar. Biol. Ecol.* 138, 1990, pp. 99-117.
- (122) Arimoto, R. and S.Y. Feng, Histological studies on mussels from dredge spoil dumpsites. *Estuar. Coastal Shelf Sci.* 17:535-546, 1983.
- (123) Devi, V. U., "Bioaccumulation and Metabolic Effects of Cadmium on Marine Fouling Dressinid Bivalve, *Mytilopsis sallei* (Recluz)," *Arch. Environ. Contam. Toxicol.* 31, 1996, pp. 47-53.
- (124) Bergen, B. J., Nelson, W. G., and Pruell, R. J., "Bioaccumulation of PCB Congeners by Blue Mussels (*Mytilus edulis*) Deployed in New Bedford Harbor, Massachusetts," *Environ. Toxicol. Chem.* 12, 1993, pp. 1671-1681
- (125) Fisher, N. S., Teysse, J. L., Fowler, S. W., and Wang, W. X., "Accumulation and Retention of Metals in Mussels from Food and Water: a Comparison of Field and Laboratory Conditions," *Environ. Sci. Technol.* 30(11), 1996, pp. 3232-3242.
- (126) Mallet, A. L. and Carver, C. E., "Comparative Growth and Survival Patterns of *Mytilus trossulus* and *Mytilus edulis* in Atlantic Canada," *Can. J. Fish. Aquat. Sci.* 52, 1995, pp. 1873-1880.
- (127) Lares, M. L. and Orians, K. J., "Natural Cd and Pb Variations in *Mytilus californianus* During the Upwelling Season," *Sci. Tot. Environ.* 197, 1997, pp. 177-195.
- (128) Kannan, N., Tanabe, S., Tatsukawa, R., and Phillips, D. J. H., "Persistency of Highly Toxic Coplanar PCB's in Aquatic Ecosystems Uptake and Release Kinetics of PCB's in Green-Lipped Mussels *Perna viridis* Linnaeus," *Environ. Pollut.* 56(1), 1989, pp. 65-76.
- (129) Wu, R. S. S., and Shin, P. K. S., "Transplant Experiments on Growth and Mortality of the Fan Mussel *Pinna bicolor*," *Aquaculture* 163(1/2), 1998, pp. 47-.
- (130) Davies, I. M., Drinkwater, J., and McKie, J. C., "Effects of Tributyltin Compounds from Antifoulants on Pacific Oysters *Crassostrea gigas* in Scottish Sea Lochs," *U. K. Aquaculture* 74(3-4), 1988, pp. 319-330.
- (131) Phelps, H. L., and Page, D. S., "Tributyltin Bioassays in Portugal with the Dogwhelk (*Nucella lapillus*) and the Portuguese Oyster (*Crassostrea angulata*)," *Mar. Env. Res.* (Submitted).
- (132) Wirth, E. F., Scott, G. I., Fulton, M. H., Van Dolah, R. F., Maier, P. P., Hadley, N., Daugomah, J. W., and Key, P. B., "In Situ Monitoring of Dredged Material Spoil Sites Using the Oyster *Crassostrea virginica*," *Arch. Environ. Contam. Toxicol.* 30, 1996, pp. 340-348.
- (133) Meyer, U., Hagen, W., and Medeiros, C., "Mercury in a Northeastern Brazilian Mangrove Area, a Case Study: Potential of the Mangrove Oyster *Crassostrea rhizophorae* as Bioindicator for Mercury," *Mar. Biol.* 131, 1998, pp. 113-121.
- (134) Cooper, R. J., Langlois, D., and Olley, J., "Heavy Metals in Tasmanian Shellfish. I. Monitoring Heavy Metal Contamination in the Derwent Estuary: Use of Oysters and Mussels," *J. Appl. Toxicol.* 2(2), 1982, pp. 99-109.
- (135) Berthou, F., Balouet, G., Bodennec, G., and Marchand, M., "The Occurrence of Hydrocarbons and Histopathological Abnormalities in Oysters for Seven Years Following the Wreck of the Amoco Cadiz in Brittany (France)," *Mar. Environ. Res.* 23, 1987, pp. 103-133.
- (136) Salazar, S. M., Salazar, M. H., Davidson, B. M., Stang, P. M., and Meyers Schulte, K. J., "A Portable Environmental Test System: A Field Assessment of Organotin Leachates," Naval Ocean Systems Center Technical Report 1202, 1987.
- (137) Scanes, P., "Uptake and Depuration of Organochlorine Compounds in Sydney Rock Oysters," *Mar. Freshwater Res.* 48, 1997, pp. 1-6.
- (138) Din, Z. B. and Ahamad, A., "Changes in the Scope for Growth of Blood Cockles (*Anadara granosa*) Exposed to Industrial Discharge," *Mar. Pollut. Bull.* 31(4-12), 1995, pp. 406-410.
- (139) Essink, K. and Bos, A. H., "Growth of Three Bivalve Mollusks Transplanted Along the Axis of the Ems Estuary, West Germany, the Netherlands," *Neth. J. Sea Res.* 19(1), 1985, pp. 45-51.
- (140) Dobbins, S. J., Barker, M. F., and Jillett, J. B., "Experimental Shore Level Transplantation of the New Zealand Cockle *Chione stutchburyi*," *J. Shellfish Res.* 8(1), 1989, pp. 197-212.
- (141) Armstrong, D. A., and Milleman, R. E., "Effects of the Insecticide Carbaryl on Clams and Some Other Intertidal Mud Flat Animals," *J. Fish. Res. Bd. Canada* 31(4), 1974, pp. 466-470.
- (142) Roesijadi, G., Anderson, J. W., and Blaylock, J. W., "Uptake of Hydrocarbons from Marine Sediments Contaminated with Prudhoe Bay Crude Oil: Influence of Feeding Type of Test Species and Availability of Polycyclic Aromatic Hydrocarbons," *J. Fish. Res. Bd. Can.* 35, 1978, pp. 608-614.
- (143) Ringwood, A. H., Van Dolah, R. F., Holland, A. F., and De Lorenzo, M. E., "Year I Demonstration Project Studies Conducted in the Carolinian Province: Results and Summaries," Charleston, South Carolina, South Carolina Department of Natural Resources, Marine Resources Research Department, 1995.
- (144) Dow, R. L., "Reduced Growth and Survival of Clams Transplanted to an Oil Spill Site," *Mar. Pollut. Bull.* 6(8), 1975, pp. 124-125.
- (145) Langston, W. J., "Availability of Arsenic to Estuarine and Marine Organisms: a Field and Laboratory Evaluation," *Mar. Biol.* 80, 1984, pp. 143-154.

- (146) Davis, C. V., Scully, K. C., and Shumway, S. E., "Juvenile and Yearling Growth of Atlantic Surfclams *Spisula solidissima* (dillwyn, 1817) in Maine," *Journal of Shellfish Research* 16(1), 1997, pp. 161-168.
- (147) Allison, E. H., and Brand, A. R., "A Mark-Recapture Experiment on Queen Scallops, *Aequipecten opercularis*, on a North Irish Sea Fishing Ground," *J. Mar. Biol. Assoc. U.K.* 75, 1995, pp. 323-335.
- (148) Rice, M. A., Rheault, R. B., Perez, M. S., and Perez, V. S., "Experimental Culture and Particle Filtration by Asian Moon Scallops, *Amusium pleuronectes*," *Asian Fisheries Sci.* 7, 1994, pp. 179-185.
- (149) Peterson, C. H., Summerson, H. C., and Luettich, R. A. Jr., "Response of Bay Scallops to Spawner Transplants: a Test of Recruitment Limitation," *Mar. Ecol. Prog. Ser.* 132, 1996, pp. 93-107.
- (150) Martinez, G., Torres, M., Uribe, E., Diaz, M. A., and Perez, H., "Biochemical Composition of Broodstock and Early Juvenile Chilean Scallops, *Argopecten purpuratus* Lamarck, Held in Two Different Environments," *J. Shellf Res.* 11(2), 1992, pp. 307-313.
- (151) Gosling, E. M., and Burnell, G. M., "Evidence for Selective Mortality in *Chlamys varia* (L.) Transplant Experiments," *J. Mar. Biol. Assoc. U. K.* 68, 1988, pp. 251-258.
- (152) MacDonald, B. A. and Bourne, N. F., "Growth of the Purple-Hinge Rock Scallop, *Crassodoma gigantea* Gray, 1825, Under Natural Conditions and Those Associated with Suspended Culture," *J. Shellfish Res.* 8(1), 1989, pp. 179-186.
- (153) Minchin, D., Duggan, C. B., and King, W., "Possible Effects of Organotins on Scallop Recruitment," *Mar. Pollut. Bull.* 18(11), 1987, pp. 604-608.
- (154) Beaumont, A. R., Morvan, C., Huelvan, S., Lucas, A., and Ansell, A. D., "Genetics of Indigenous and Transplanted Populations of *Pecten maximus*: No Evidence for the Existence of Separate Stocks," *J. Exp. Mar. Biol. Ecol.* 169, 1993, pp. 77-88.
- (155) Cranford, P. J. and Hargrave, B. T., "In Situ Time-Series Measurement of Ingestion and Absorption Rates of Suspension-Feeding Bivalves: *Placopecten magellanicus*," *Limnol. Oceanogr.* 39(3), 1994, pp. 730-738.
- (156) Sheehan, R. J., Neves, R. J., and Kitchel, H. E., "Fate of Freshwater Mussels Transplanted to Formerly Polluted Reaches of the Clinch and North Fork Holston Rivers, Virginia," *J. Freshw. Ecol.* 5(2), 1992, pp. 139-149.
- (157) Waller, D. L., Rach, J. J., Cope, W. G., and Miller, G. A., "Effects of Handling and Aerial Exposure on the Survival of Unionid Mussels," *J. Fresh. Ecol.* 10(3), 1995, pp. 199-207.
- (158) Adams, T. G., Atchison, G. J., and Vetter, R. J., "The Use of the Three-Ridge Clam (*Amblema perplicata*) to Monitor Trace Metal Contamination," *Hydrobiologia* 83, 1981, pp. 67-72.
- (159) Englund, V.P.M. and M.P. Heino, Valve movement of the freshwater mussel *Anodonta anatina*: a reciprocal transplant experiment between lake and river. *Hydrobiologia* 328(1):49
- (160) Hayer, F., Wagner, P., and Phan, J. C., "Monitoring of Extractable Organic Halogens (EOX) in Chlorine Bleached Pulp and Paper Mill Effluents Using Four Species of Transplanted Aquatic Mollusks," *Chemosphere* 33(11), 1996, pp. 2321-2334.
- (161) Hanson, J. M., Mackay, W. C., and Prepas, E. E., "The Effects of Water Depth and Density on the Growth of a Unionid Clam," *Freshwat. Biol.* 19, 1988, pp. 345-355.
- (162) Rutherford, L. A., Ernst, W. R., Doe, K. G., and Hennigar, P. A., "Chemical Characterization, Aquatic Toxicity, and Environmental Impact of Untreated Effluent Discharges from Three Textile Mills in the Atlantic Region," E. G. Baddaloo, S. Ramamoorthy, and J. W. Moore, eds., *Proceedings of the Nineteenth Annual Aquatic Toxicity Workshop* : October 4-7, 1992, Edmonton, Alberta. Canadian Technical Report of Fisheries and Aquatic Sciences, 1992, pp. 293-299.
- (163) Lopes, J. L. C., Casanova, I. C., de Figueireido, M. C. G., Nather, F. C., and Avelar, W. E. P., "*Anodontites trapesialis*: A Biological Monitor of Organochlorine Pesticides," *Arch. Environ. Contam. Toxicol.* 23, 1992, pp. 351-354.
- (164) Kraak, M. H. S., Scholten, M. C. Th., Peeters, W. H. M., and de Kock, W. C., "Biomonitoring of Heavy Metals in the Western European Rivers Rhine and Meuse Using the Freshwater Mussel *Dreissena polymorpha*," *Environ. Pollut.* 74, 1991, pp. 101-114.
- (165) Kauss, P. B., and Hamdy, Y. S., "Biological Monitoring of Organochlorine Contaminants in the St. Clair and Detroit Rivers Using Introduced Clams, *Elliptio complanata*," *J. Great Lakes Res.* 11(3), 1985, pp. 247-263.
- (166) Trdan, R. J., and Hoeh, W. R., "Relocation of Two State-Listed Freshwater Mussel Species (*Epioblasma torulosa rangiana* and *Epioblasma triquetra*) in Michigan," K. S. Cummings, A. C. Buchanan, and L. M. Koch, eds., *Conservation and Management of Freshwater Mussels, Proceedings of a UMRCC Symposium*, 12-14 October 1992, St. Louis, MO. Rock Island, Upper Mississippi River Conservation Committee. 1993, pp. 100-105.
- (167) Burggraaf, S., Langdon, A. G., Wilkins, A. L., and Roper, D. S., "Accumulation and Depuration of Resin Acids and Fichtelilte by the Freshwater Mussel *Hyridella menziesi*," *Environ. Toxicol. Chem.* 15(3), 1996, pp. 369-375.
- (168) Nelson, D., "Relocation of *Lampsilis higginsii* in the Upper Mississippi River," A. C. Miller, ed., *Report of Freshwater Mussels Workshop*, St. Louis, Missouri, 26-27, October 1982. Vicksburg, US Army Engineer Waterways Experiment Station. 1983, pp. 104-107.
- (169) Hinch, S. G., Bailey, R. C., and Green, R. H., "Growth of *Lampsilis radiata* (Bivalvia: Unionidae) in Sand and Mud: a Reciprocal Transplant Experiment," *Can. J. Fish. Aquat. Sci.* 43, 1986, pp. 548-552.
- (170) Czarnecki, J. M., "Use of the Pocketbook Mussel *Lampsilis ventricosa* for Monitoring Heavy Metal Pollution in an Ozark Stream," *Bull. Environ. Contam. Toxicol.* 38, 1987, pp. 641-646.
- (171) Hubbs, D., Hunt, T., and Kathman, R. D., "Lemiox Rimosus Translant Site Survey," Report prepared for the Upper Duck River Development Agency. Young-Morgan & Associates, Inc., Franklin, 1991, 26 pp.
- (172) Fishman, P. A., and Johnson, S. R., Quartz Mountain Gold Project Clam Bioaccumulation Study. Prepared for Galactic Services, Inc., 1989.
- (173) Sparks, R. E. and Blodgett, K. D., "Effects of Three Commercial Harvesting Methods on Mussel Beds," Ill. Nat. Hist. Survey, Aquat. Biol. Sect. Tech. Rept. (10), 1983, 44 pp.
- (174) Harris, J. L., "Relocation of the Fat Pocketbook Pearly Mussel, *Proptera Capax* (Green), in the St. Francis River at Madison, St. Francis County, Arkansas," Environmental Division, Arkansas State Highway and Transportation Department, Little Rock, 1986, 14 pp.
- (175) Foster, R. B., and Bates, J. M., "Use of Freshwater Mussels to Monitor Point Source Industrial Discharges," *Environ. Sci. Tech.* 12, 1978, pp. 958-962.
- (176) Storey, A. W. and Edward, D. H. D., "The Freshwater Mussel, *Westraulino carteri* Iredale, as a Biological Monitor of Organochlorine Pesticides," *Aust. J. Mar. Freshwat. Res.* 40, 1989, pp. 587-593.
- (177) Scanes, P., "Trace Metal Uptake in Cockles *Anadara trapezium* from Lake Macquarie, New South Wales," *Mar. Ecol. Prog. Ser.* 102, 1993, pp. 135-142.
- (178) Doherty, F.G. The Asiatic clam, *Corbicula* spp., as a biological monitor in freshwater environments. *Environ. Monit. Assess.* 15:143-181, 1990.

- (179) Ohyama, T., Jin, K., Katoh, Y., Chiba, Y., and Inoue, K., "1,3,5-Trichloro-2- (4- Nitrophenoxy)-Benzene (CNP) in Water, Sediments, and Shellfish of the Ishikari River," *Bull. Environ. Contam. Toxicol.* 37(3), 1986, pp. 344-349.
- (180) Hartley, D. M., and Johnston, J. B., "Use of the Freshwater Clam *Corbicula manilensis* as a Monitor for Organochlorine Pesticides," *Bull. Environ. Contam. Toxicol.* 31, 1983, pp. 33-40.
- (181) Zischke, J. A., and Arthur, J. W., "Effects of Elevated Ammonia Levels on the Fingernail Clam, *Musculium transversum* , in Outdoor Experimental Streams," *Arch. Environ. Contam. Toxicol.* 16, 1987, pp. 336-231.
- (182) Deleon, I. R., Ferrario, J. B., and Byrne, C. J., "Bioaccumulation of Polynuclear Aromatic Hydrocarbons by the Clam, *Rangia cuneata*, in the Vicinity of a Creosote Spill," *Bull. Env. Contam. Toxicol.*, Vol 41, 1988, pp. 872-879.
- (183) Rice, C. P., and White, D. S., "PCB Availability Assessment of River Dredging Using Caged Clams and Fish," *Environ. Toxicol. Chem.* 6, 1987, pp. 259-274.
- (184) Applied Biomonitoring, "Progress Report #1, City of Winnipeg: Caged Bivalve Study," Prepared for TetrES, 1999, July 23, 1999.
- (185) Belanger, S. E., Cherry, D. S., and Cairns, J., "Uptake of Chrysotile Asbestos Fibers Alters Growth and Reproduction of Asiatic Clams.," *Can. J. Fish. Aquat. Sci.*, Vol 43, 1986, pp. 43-52.
- (186) Belanger, S. E., Farris, J. L., Cherry, D. S., and Cairns Jr., "Growth of Asiatic Clams (*Corbicula* sp.) During and After Long-Term Zinc Exposure in Field-Located and Laboratory Artificial Streams," *Arch. Environ. Contam. Toxicol.*, Vol 15, 1986, pp. 427-434.
- (187) Belanger, S. E., Cherry, D. S., Cairns, J., and McGuire, M. J., "Using Asiatic Clams As a Biomonitor for Chrysotile Asbestos in Public Water Supplies," *Journal AWWA*, March 1987, 1987, pp. 69-74.
- (188) Belanger, S. E., Farris, J. L., Cherry, D. S., and Cairns, J., "Validation of *Corbicula fluminea* Growth Reductions Induced by Copper in Artificial Streams and River Systems," *Can. J. Fish. Aquat. Sci.* Vol 47, 1990, pp. 904-914.
- (189) Malley, D. F., Stewart, A. R., and Hall, B. D., "Uptake of Methyl Mercury by the Floater Mussel, *Pyganodon grandis* (Bivalvia, Unionidae) Caged in a Flooded Wetland," *Environ. Toxicol. Chem.*, Vol 15, No. 6, 1996, pp. 928-936.
- (190) US Department of Interior Fish and Wildlife Service, "Common Rangia. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Gulf of Mexico)," *Biological Report 82 (11.31) - TR EL- 82-4*, US Army Corps of Engineers Waterways Experiment Station, 1985.
- (191) Demas, C. R., and Demcheck, D. K., "Uptake of Manmade Organic Compounds by *Rangia cuneata* in the Lower Calcasieu River, Louisiana," *U.S. Geological Survey Toxic Substances Hydrology Program - Surface-Water Contamination: Proceedings of the Technical Meeting*, Denver, CO, 1987.
- (192) Fucik, K. W., Armstrong, H. W., and Neff, J. M., "Uptake of Naphthalenes by the Clam *Rangia cuneata*, in the Vicinity of an Oil Separator Platform in Trinity Bay, Texas," *Proceedings, 1977 Oil Spill Conference (Prevention, Behavior, Control, Cleanup)*. March 8-10, 1977, New Orleans, LA, pp. 637-640.
- (193) Lunsford, C. A., and Blem, C. R., "Annual Cycle of Kepone Residue and Lipid Content of the Estuarine Clam, *Rangia cuneata*," *Estuaries*, Vol 5, No. 2, 1995, pp. 121-130.
- (194) Kennedy, V. S., Newell, R. I. E., and Eble, A. F., *The eastern oyster, Crassostrea virginica*, Maryland Sea Grant, College Park, MD. 1996.
- (195) Waldock, M. J., Thain, J. E., and Waite, M. E., "An Assessment of the Value of Shell Thickening in *Crassostrea gigas* As an Indicator of Exposure to Tributyltin," *Tributyltin: Environmental Fate and Effects*, M. A. Champ, and P. F. Seligman, eds., Chapman and Hall, London, 1996, pp. 219-237.
- (196) Granmo, A., "Mussels As a Tool in Impact Assessment," *Phuket Mar. Biol. Cent. Spec. Publ. No. 15*, 1995, pp. 215-220.
- (197) Milam, C. D., Farris, J. L., Dwyer, J. F., and Hardesty, D. K., "Comparison of Acute Toxicity Responses of *Glochidia* (Unionidae) to Six Chemicals (Abstract Only)," *SETAC 19th Annual Meeting. The Natural Connection: Environmental Integrity and Human Health*, Charlotte, NC, Society of Environmental Toxicology and Chemistry, 1998.
- (198) Beaumont, A. R., Tserpes, G., and Budd, M. D., "Some Effects of Copper on the Veliger Larvae of the Mussel *Mytilus edulis* and the Scallop *Pecten maximum* (Molluscs, Bivalvia)," *Mar. Environ. Res.* Vol 21, 1987, pp. 299-309.
- (199) Butler, R., Roddie, B. D., and Mainstone, C. P., "The Effects of Sewage Sludge on Two Life-History Stages of *Mytilus edulis*," *Chemistry and Ecology*, Vol 4, 1990, pp. 211-219.
- (200) Gardner, J. P., "The *Mytilus edulis* Species Complex in Southwest England: Effects of Hybridization and Introgression Upon Interlocus Associations and Morphometric Variation," *Mar. Biol.*, Vol 125, 1996, pp. 385-399.
- (201) Kenchington, E., Landry, D., and Bird, C. J., "Comparison of Taxa of the Mussel *Mytilus* (Bivalvia) by Analysis of the Nuclear Small-Subunit rRNA Gene Sequence," *Can. J. Fish. Aquat. Sci.*, Vol 52, 1995, pp. 2613-2620.
- (202) Suchanek, T. H., Geller, J. B., Kreiser, B. R., and Mitton, J. B., (Submitted), "Distributions of the Sibling Species *Mytilus Galloprovincialis* and *M. Trossulus* (Bivalvia: Mytilidae) and Their Hybrids in the North Pacific," *Biol. Bull.*
- (203) McDonald, J. H., and Koehn, R. K., "The Mussels *Mytilus galloprovincialis* and *M. trossulus* on the Pacific Coast of North America," *Mar. Biol.*, Vol 99, 1988, pp. 111-118.
- (204) Bates, J. A., and Innes, D. J., "Genetic Variation Among Populations of *Mytilus* spp. in Eastern Newfoundland," *Mar. Biol.*, Vol 124, 1995, pp. 417-424.
- (205) Gunther, A. J., Davis, J. A., Hardin, D. D., Gold, J., Bell, D., Crick, J. R., Scelfo, G. M., Sericano, J., and Stephenson, M., "Long-Term Bioaccumulation Monitoring with Transplanted Bivalves in the San Francisco Estuary," *Mar. Pollut. Bull.*, Vol 38, No. 3, 1999, pp. 170-181.
- (206) Rand, G.M. *Fundamentals of Aquatic Toxicology: Effects, Environmental Fate, and Risk Assessment*. North Plam Beach, Florida, Taylor and Francis. 1995.

ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

*This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or [service@astm.org](mailto:service@astm.org) (e-mail); or through the ASTM website ([www.astm.org](http://www.astm.org)).*